



Billing and coding guide

For questions or support, reach out to your Novartis Access and Reimbursement Expert

Name: _____ Phone: _____

Title: _____ Email: _____

You can also talk to a dedicated Access Specialist at the LEQVIO[®] Service Center



Phone: **833-LEQVIO2 (833-537-8462)**



Fax: **877-LEQVIO8 (877-537-8468)**



Website: **LEQVIOhcp.com**



Portal: **ServiceCenterPortal.com**

Overview

This guide is intended to provide an overview of coding and coverage information related to LEQVIO®. Health care professionals can reference this guide, in addition to other sources of information, to determine for themselves the appropriate claims to file for LEQVIO and the related services. Novartis does not guarantee payment or coverage for any product or service.

The health care billing environment is constantly evolving to keep pace with scientific advances and financial constraints. Information specific to billing and coding is subject to change without notice and should be verified by the provider for each patient prior to treatment. A provider should contact the patients' payers directly for any revised or additional requirements, information, or guidance.

It is always the provider's responsibility to determine the appropriate health care setting, and to submit true and correct claims for the products and services rendered. Providers should contact third-party payers for specific information on their coding, coverage, payment policies, and fee schedules.

The site of care determines specific coding requirements. It is always the provider's responsibility to determine medical necessity for a specific service, the proper site for delivery of any services, and to submit appropriate codes, charges, and modifiers for services that are rendered.

"Patient" in this document may refer to adult patients or a pediatric or adult patient's authorized representative.

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

Basic coverage information

Billing and coding requirements for LEQVIO® will vary based on many factors, including the site of service where the drug is administered, the type of insurance the patient has, and the benefit under which LEQVIO is covered.

Site of service

LEQVIO may be administered in physicians' offices or in hospital outpatient departments. For most payers, the site of service will affect the billing and coding requirements. This guide provides information on coverage, coding, and billing for LEQVIO when administered in physicians' offices, hospital outpatient settings, and stand-alone alternate sites of care.

Payer type

Coverage, as defined by each payer type and benefit package, may vary depending on the site of service and the patient's status and medical history.



Medicare

Medicare typically covers and separately reimburses drugs provided in the physician's office as well as most drugs provided in the hospital outpatient department that are not self-administered and are provided incident to a physician service.

Coverage under Medicare Advantage can vary by plan. Providers should check with the patient's plan for specific coverage and payment information



Private payers

Private payers may cover LEQVIO and the medical services associated with its administration. However, there may be restrictions on coverage, such as special requirements for distribution and precertification. Private payers may also vary in the payment methods they use to reimburse the sites of service where LEQVIO is administered



Medicaid

Medicaid coverage and payment for LEQVIO can vary by state or by the specific managed Medicaid plan. Providers should check with the state program or plan for specific coverage information and all payer types for fee schedules

Benefit category

Most payers cover physician-administered products such as LEQVIO under a medical benefit rather than a pharmacy benefit. In the case of Medicare, LEQVIO will typically be covered under Part B. However, private payers and Medicaid may require that physicians obtain LEQVIO through a specialty pharmacy. Specialty pharmacies may bill the payer under the medical or pharmacy benefit, depending on what that payer requires. Additionally, Medicare Advantage payers may cover LEQVIO in a similar way to private payers depending on the benefit design of the plan.

Reference: Tikkanen R, et al. Published June 5, 2020. Accessed August 11, 2025. <https://www.commonwealthfund.org/international-health-policy-center/countries/united-states>

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

Relevant codes: drug codes

National Drug Code (NDC)¹

The NDC is a product identifier for drugs in the United States present on all over the counter and prescription medication packages and inserts. The NDC number is essential for proper claim processing when submitting claims for drugs used. As not all NDCs are set up the same, the table below demonstrates how to achieve the 11-digit NDC for LEQVIO®.

NOTE: Be sure that hyphens are excluded from submission on the claim and check with the patient's health insurance provider to determine sequence requirements.

Tradename	Package strength	10-digit	NDC number	11-digit	NDC number for payer
LEQVIO	284 mg/1.5 mL single-dose prefilled syringe	4-4-2	0078-1000-60	5-4-2	00078-1000-60

Healthcare Common Procedure Coding System (HCPCS) level II code²

HCPCS Level II codes are used to identify drugs, supplies, medical procedures, and other services.

HCPCS code	Descriptor	Billing units
J1306	Injection, inclisiran, 1 mg	284

Modifiers^{3,4}

Modifiers provide additional information about a service or procedure without changing its definition or code. By using modifiers, health care providers can indicate specific circumstances such as route of administration, wasted product, and more. To determine the applicability of a modifier, it is recommended to consult the relevant CMS manuals.

NOTE: the modifiers below will accompany the HCPCS code within the claim form.

Modifier	Information
JZ	Providers and suppliers are required to report the JZ modifier on all claims that bill for drugs from single-dose containers that are separately payable under Medicare Part B when there are no discarded amounts
TB	340B covered entities to only report the TB modifier on claims for products acquired through the 340B program. This update impacts Medicare providers and suppliers who bill for separately payable Part B drugs and biologics and participate in the 340B Pricing Program

References: **1.** Leqvio. Prescribing information. Novartis Pharmaceuticals Corp. **2.** Centers for Medicare & Medicaid Services. CMS HCPCS Application Summaries and Coding Recommendations: First Quarter, 2022 HCPCS Coding Cycle. Accessed August 11, 2025. <https://www.cms.gov/files/document/2022-hcpcs-application-summary-quarter-1-2022-drugs-and-biologicals.pdf> **3.** Centers for Medicare & Medicaid Services. Medicare program: discarded drugs and biologics—JW modifier and JZ modifier policy frequently asked questions. Accessed August 11, 2025. <https://www.cms.gov/medicare/medicare-fee-for-service-payment/hospitaloutpatientpps/downloads/jw-modifier-faqs.pdf> **4.** Centers for Medicare & Medicaid Services. Revised Part B inflation rebate guidance: Use of the 340B modifier. Published December 14, 2023. Accessed August 11, 2025. <https://www.cms.gov/files/document/revised-part-b-inflation-rebate-340b-modifier-guidance.pdf>

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

Relevant codes: CPT and setting of care codes

Current Procedural Terminology (CPT) code¹

CPT codes are the most widely accepted codes for reporting medical procedures and services under government and private health plans.

CPT code*	Description
96372	Therapeutic, Prophylactic, and Diagnostic Injection (specific substance or drug; subcutaneous or intramuscular)

Place of service (POS) codes²

POS codes are used to indicate the setting in which a service was provided. CMS maintains a database of POS codes commonly used in the health care industry. Below are examples of some common POS codes. Review the full listing of the POS codes on the CMS website and consult your health plan's guidance to determine the correct code for your institution.

POS code	Description
11	Office: Location, other than a hospital, skilled nursing facility (SNF), military treatment facility, community health center, state or local public health clinic, or intermediate care facility (ICF), where the health professional routinely provides health examinations, diagnosis, and treatment of illness or injury on an ambulatory basis
22	On Campus—Outpatient Hospital: A portion of a hospital's main campus which provides diagnostic, therapeutic (both surgical and nonsurgical), and rehabilitation services to sick or injured persons who do not require hospitalization or institutionalization

Revenue codes³

Specific forms, such as the UB-04 (CMS-1450) form, require documentation of revenue codes associated with services provided to patients.

Revenue code	Description
0636	Drugs requiring detailed coding
0250	General pharmacy
0500	General outpatient services
0510	General clinic services

*CPT © 2026 American Medical Association. All rights reserved.

References: **1.** AAPC Coder. Accessed August 11, 2025. <https://www.aapc.com/codes/cpt-codes/96372> **2.** Centers for Medicare & Medicaid Services. Place of Service Codes for Professional Claims. Accessed August 11, 2025. <https://www.cms.gov/medicare/medicare-fee-for-service-payment/physicianfeesched/downloads/website-pos-database.pdf> **3.** Noridian Healthcare Solutions. Revenue Codes. Accessed August 11, 2025. <https://med.noridianmedicare.com/web/jea/topics/claim-submission/revenue-codes>

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

Physician's office: sample CMS-1500 claim form

LEQVIO® and the associated services provided in a physician office are billed on the CMS-1500 claim form or its electronic equivalent. A sample CMS-1500 claim form for billing LEQVIO is provided below.¹

The sample claim form provided below is only an example. It is always the provider's responsibility to determine the appropriate health care setting and to submit true and correct claims for the products and services rendered. Providers should contact third-party payers for specific information on their coding, coverage, payment policies, and fee schedules.

- Box 21**
Relevant diagnosis code(s) (ICD-10-CM).
- Box 23**
Prior authorization number, if available.
- Box 24A**
In the non-shaded area, list the date of service. In the shaded area, give a detailed drug description. List the N4 indicator first, then the 11-digit NDC number. Third is the unit of measurement qualifier; the unit quantity is listed at the end.
Example N400078100060ML15
- Box 24B**
Enter the appropriate POS code to indicate the setting where a service was provided.
- Box 24D**
Enter the appropriate HCPCS code J1306 for LEQVIO use as required by the payer.² The HCPCS code must be accompanied by the JZ modifier, indicating zero drug wasted.³ 340B covered entities must use the TB modifier accompanied by the HCPCS code.⁴ Include the appropriate CPT code to report the administration procedure, 96372.⁵
- Box 24E**
Enter the diagnosis code reference letter (A or B) as shown in Box 21 to relate the date of service and the procedures performed to the primary diagnosis. If there is more than one diagnosis required for a procedure code, only reference one letter from Box 21.
- Box 24G**
Include the appropriate number of billing units for LEQVIO: 284 mg=284 billing units.

IMPORTANT INFORMATION: The coding, coverage, and payment information contained herein is gathered from various resources, general in nature, and subject to change without notice. Third-party payment for medical products and services is affected by numerous factors. It is always the provider's responsibility to determine the appropriate health care setting and to submit true and correct claims for those products and services rendered. Providers should contact third-party payers for specific information on their coding, coverage, and payment policies. Information and materials are provided to assist health care providers, but the responsibility to determine coverage, reimbursement, and appropriate coding for a particular patient and/or procedure remains, at all times, with the provider.

References: **1.** Centers for Medicare & Medicaid Services. Accessed August 11, 2025. <https://www.cms.gov/Medicare/CMS-Forms/CMS-Forms/Downloads/CMS1500.pdf> **2.** Centers for Medicare & Medicaid Services. CMS HCPCS Application Summaries and Coding Recommendations: First Quarter, 2022 HCPCS Coding Cycle. Accessed August 11, 2025. <https://www.cms.gov/files/document/2022-hcpcs-application-summary-quarter-1-2022-drugs-and-biologicals.pdf> **3.** Centers for Medicare & Medicaid Services. Medicare program: discarded drugs and biologicals—JW modifier and JZ modifier policy frequently asked questions. Accessed August 11, 2025. <https://www.cms.gov/medicare/medicare-fee-for-service-payment/hospitaloutpatientpps/downloads/jw-modifier-faqs.pdf> **4.** Medicare Part B inflation rebate guidance: use of the 340b modifier. Medicare Learning Network Fact Sheet. Published December 2023. Accessed August 11, 2025. <https://www.cms.gov/files/document/mln4800856-medicare-part-b-inflation-rebate-guidance-use-340b-modifier.pdf> **5.** AAPC Coder. Accessed August 11, 2025. <https://www.aapc.com/codes/cpt-codes/96372>

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.



Hospital outpatient: sample CMS-1450 (UB-04) form

LEQVIO® and the associated services provided in a hospital outpatient setting are billed on the UB-04 claim form or its electronic equivalent. A sample UB-04 claim form for billing LEQVIO is provided below.¹

The sample claim form provided below is only an example. It is always the provider's responsibility to determine the appropriate health care setting and to submit true and correct claims for the products and services rendered. Providers should contact third-party payers for specific information on their coding, coverage, payment policies, and fee schedules.

The form is a CMS-1450 (UB-04) claim form. Key sections include:

- Box 42:** Statement covers period from 01/01/2022 to 12/31/2022.
- Box 43:** Drug description: N400078100060ML1.5 Outpatient clinic. HCPCS code: J1306 JZ TB 96372. Serv units: 284.
- Box 44:** HCPCS code J1306 for LEQVIO use.
- Box 46:** Treatment authorization codes: 01, 02, 03, 04, 05, 06, 07, 08, 09, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 00.
- Box 63:** Treatment authorization codes: 01, 02, 03, 04, 05, 06, 07, 08, 09, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 00.
- Box 66:** Relevant diagnosis code(s) (ICD-10-CM): 01, 02, 03, 04, 05, 06, 07, 08, 09, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 00.

Box 42
Enter the appropriate revenue code corresponding with the HCPCS code in Box 44 (eg, 0636 revenue code for drugs requiring detailed coding, and/or 0250 for general pharmacy). Then enter the appropriate revenue code corresponding with the CPT code in Box 44 (eg, 0500 for general outpatient services or 0510 for general clinic services).²

Box 43
Enter a detailed drug description for the payer. The N4 indicator is listed first, the 11-digit National Drug Code number is listed second, a code describing the unit of measurement qualifier is listed third (eg, mL for milliliters), and the unit quantity is listed at the end. Example N400078100060ML1.5

Box 44
Enter the appropriate HCPCS code J1306 for LEQVIO use as required by the payer.³ The HCPCS code must be accompanied by the JZ modifier, indicating zero drug wasted.⁴ 340B covered entities must use the TB modifier accompanied by the HCPCS code.⁵ To report the administration procedure, enter an appropriate CPT code, 96372.⁶

Box 46
Include the appropriate number of billing units for LEQVIO: 284 mg=284 billing units. Some payers may refer to the actual quantity administered via Box 43.

Box 63
Treatment authorization codes.

Box 66
Relevant diagnosis code(s) (ICD-10-CM).

IMPORTANT INFORMATION: The coding, coverage, and payment information contained herein is gathered from various resources, general in nature, and subject to change without notice. Third-party payment for medical products and services is affected by numerous factors. It is always the provider's responsibility to determine the appropriate health care setting and to submit true and correct claims for those products and services rendered. Providers should contact third-party payers for specific information on their coding, coverage, and payment policies. Information and materials are provided to assist health care providers, but the responsibility to determine coverage, reimbursement, and appropriate coding for a particular patient and/or procedure remains, at all times, with the provider.

References: **1.** Centers for Medicare & Medicaid Services. Accessed August 11, 2025. <https://www.cms.gov/Regulations-and-Guidance/Legislation/PaperworkReductionActof1995/PRA-Listing-Items/CMS-1450> **2.** Noridian Healthcare Solutions. Revenue Codes. Accessed August 11, 2025. <https://med.noridianmedicare.com/web/jea/topics/claim-submission/revenue-codes> **3.** Centers for Medicare & Medicaid Services. CMS HCPCS Application Summaries and Coding Recommendations: First Quarter, 2022 HCPCS Coding Cycle. Accessed August 11, 2025. <https://www.cms.gov/files/document/2022-hcpcs-application-summary-quarter-1-2022-drugs-and-biologicals.pdf> **4.** Centers for Medicare & Medicaid Services. Medicare program: discarded drugs and biologicals–JW modifier and JZ modifier policy frequently asked questions. Accessed August 11, 2025. <https://www.cms.gov/medicare/medicare-fee-for-service-payment/hospitaloutpatientpps/downloads/jw-modifier-faqs.pdf> **5.** Medicare Part B inflation rebate guidance: use of the 340b modifier. Medicare Learning Network Fact Sheet. Published December 2023. Accessed August 11, 2025. <https://www.cms.gov/files/document/mln4800856-medicare-part-b-inflation-rebate-guidance-use-340b-modifier.pdf> **6.** AAPC Coder. Accessed August 11, 2025. <https://www.aapc.com/codes/cpt-codes/96372>



Claim filing checklist

Once your patient is administered LEQVIO®, a claim for reimbursement is submitted to their health insurance provider. When submitting a claim, it is a best practice to understand the payer's specific billing and coding requirements and ensure that the submitted claim is as complete as possible. You may want to reference the following general tips when filing claims for LEQVIO:

- Use appropriate codes to report the patient's condition, the drugs the patient received, and the services you have provided**
 - ICD-10-CM code
 - For a list of potentially applicable diagnosis codes, please see the [ICD-10-CM codes tab](#)
 - NDC
 - CPT code
 - HCPCS code
 - JZ modifier
 - 340B modifier

- Attach additional information to the claim if necessary**
 - Letter of medical necessity
 - Prescribing Information
 - Patient notes

- Review claim for accuracy, including patient identification numbers and coding**

- File claim as soon as possible and within health plan filing time limits**

- Reconcile claim reports promptly and thoroughly to ensure claims have been appropriately processed and paid**

- Verify that payment amounts correspond with your public health plan allowables and your private health plan contracts**

For a list of potentially applicable diagnosis codes, please see the [ICD-10-CM codes tab](#)

Novartis Patient Support is here to help

Enroll in Novartis Patient Support services with your patients to receive:



ACCESS & REIMBURSEMENT SUPPORT

A dedicated Access Specialist can provide you and your office with insurance, acquisition, and reimbursement support.



PATIENT AFFORDABILITY SUPPORT

The LEQVIO® Co-pay Program can help eligible commercially insured patients **pay as little as \$0** for LEQVIO.

Subject to terms and conditions. Limitations apply.*



PATIENT EDUCATION AND ONGOING SUPPORT

The LEQVIO Care Program offers dedicated one-on-one support that can help your patients start and stay on their LEQVIO treatment.



GET STARTED

Simply download the one-page **Start Form** and fax it to **877-LEQVIO8 (877-537-8468)**

OR

Register for the Service Center Portal and submit your request online at **ServiceCenterPortal.com**

Have questions? We are here to help.



Phone: **833-LEQVIO2 (833-537-8462)**



Fax: **877-LEQVIO8 (877-537-8468)**



Website: **LEQVIOhcp.com**



Portal: **ServiceCenterPortal.com**

***Limitations apply.** Valid only for those with commercial insurance. The Program may include the Co-pay Card, Payment Card (if applicable), and Rebate. Per-treatment maximums and an annual benefit cap apply. For patients covered under the medical benefit, rebate for out-of-pocket costs will be assigned directly to provider, unless patient requests direct reimbursement. Patient is responsible for any costs once limit is reached in a calendar year. Program not valid (i) under Medicare, Medicaid, TRICARE, VA, DoD, or any other federal or state health care program, (ii) where patient is not using insurance coverage at all, (iii) where the patient's insurance plan reimburses for the entire cost of the drug, or (iv) where product is not covered by patient's insurance. The value of this program is exclusively for the benefit of patients and is intended to be credited towards patient out-of-pocket obligations and maximums, including applicable co-payments, coinsurance, and deductibles. Program is not valid where prohibited by law. Patient may not seek reimbursement for the value received from this program from other parties, including any health insurance program or plan, flexible spending account, or health care savings account. Patient is responsible for complying with any applicable limitations and requirements of their health plan related to the use of the Program. Valid only in the United States and Puerto Rico. This Program is not health insurance. Program may not be combined with any third-party rebate, coupon, or offer. Proof of purchase may be required. Novartis reserves the right to rescind, revoke, or amend the Program and discontinue support at any time without notice.

*LEQVIO Care Program is a patient support program and NOT intended to take the place of the care provided by doctors or their office staff. LEQVIO Care does not provide medical advice or treatment.

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

Potential ICD-10-CM codes

The codes listed in this tab are provided for educational purposes only and are not a guarantee of coverage or reimbursement. Coverage and reimbursement may vary significantly by payer, plan, patient, and setting of care. It is the sole responsibility of the health care provider to select the proper codes and ensure the accuracy of all statements used in seeking coverage and reimbursement for an individual patient. The codes on the following pages are included as examples of potential codes that may be relevant for LEQVIO®.

Select a link from below to view the specific codes in each category.

Primary diagnosis codes: adult patients (aged 18+ years)

Hypercholesterolemia

- [Disorders of lipoprotein metabolism and other lipidemias](#) 11

Heterozygous familial hypercholesterolemia

- [Familial hypercholesterolemia](#) 11

Primary diagnosis codes: pediatric patients (aged 12-17 years)

Heterozygous and homozygous familial hypercholesterolemia

- [Familial hypercholesterolemia](#) 11

Secondary diagnosis codes: adult patients (aged 18+ years)

Clinical ASCVD

- [Atherosclerosis and atherosclerotic heart disease](#) 11
- [Ischemic heart disease \(other\)](#) 13
- [ST elevation \(STEMI\) and non-ST elevation \(NSTEMI\) myocardial infarction](#) 14
- [Presence of cardiac and vascular implants and grafts and other postprocedural states](#) 14
- [Occlusion and stenosis of precerebral and cerebral arteries, not resulting in cerebral infarction](#) 14
- [Cerebrovascular diseases \(other\)](#) 15
- [Cerebral infarction](#) 16
- [Transient cerebral ischemic attack](#) 19
- [Vascular syndromes of brain in cerebrovascular diseases](#) 19
- [Atherosclerosis](#) 19
- [Diseases of arteries, arterioles, and capillaries \(other\)](#) 30
- [Arterial embolism and thrombosis](#) 30
- [Atheroembolism](#) 31
- [Peripheral vascular angioplasty](#) 31
- [Disorders of sphingolipid metabolism and other lipid storage disorders](#) 31

Increased risk of ASCVD

- [Type 2 diabetes mellitus](#) 31
- [Chronic kidney disease](#) 31
- [Essential \(primary\) hypertension](#) 31
- [Family history](#) 31

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

International Classification of Diseases, 10th Revision, Clinical Modification (ICD-10-CM) diagnosis codes

Primary diagnosis codes: adult patients (aged 18+ years)

Hypercholesterolemia

ICD-10-CM diagnosis code	Description
Disorders of lipoprotein metabolism and other lipidemias	
E78.00	Pure hypercholesterolemia, unspecified
E78.2	Mixed hyperlipidemia
E78.49	Other hyperlipidemia, familial combined hyperlipidemia
E78.5	Hyperlipidemia, unspecified

Heterozygous familial hypercholesterolemia

Familial hypercholesterolemia	
E78.011	Familial hypercholesterolemia, heterozygous (HeFH)
E78.019	Familial hypercholesterolemia (unspecified)

Primary diagnosis codes: pediatric patients (aged 12 to 17 years)

Heterozygous and homozygous familial hypercholesterolemia

Familial hypercholesterolemia	
E78.011	Familial hypercholesterolemia, heterozygous (HeFH)
E78.010	Familial hypercholesterolemia, homozygous (HoFH)

Secondary diagnosis codes: adult patients (aged 18+ years)

Clinical ASCVD codes

Atherosclerosis and atherosclerotic heart disease	
I25.10	Atherosclerotic heart disease of native coronary artery without angina pectoris
I25.110	Atherosclerotic heart disease of native coronary artery with unstable angina pectoris
I25.111	Atherosclerotic heart disease of native coronary artery with angina pectoris with documented spasm
I25.118	Atherosclerotic heart disease of native coronary artery with other forms of angina pectoris
I25.119	Atherosclerotic heart disease of native coronary artery with unspecified angina pectoris

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I25.700	Atherosclerosis of coronary artery bypass graft(s), unspecified, with unstable angina pectoris
I25.701	Atherosclerosis of coronary artery bypass graft(s), unspecified, with angina pectoris with documented spasm
I25.708	Atherosclerosis of coronary artery bypass graft(s), unspecified, with other forms of angina pectoris
I25.709	Atherosclerosis of coronary artery bypass graft(s), unspecified, with unspecified angina pectoris
I25.710	Atherosclerosis of autologous vein coronary artery bypass graft(s) with unstable angina pectoris
I25.711	Atherosclerosis of autologous vein coronary artery bypass graft(s) with angina pectoris with documented spasm
I25.718	Atherosclerosis of autologous vein coronary artery bypass graft(s) with other forms of angina pectoris
I25.719	Atherosclerosis of autologous vein coronary artery bypass graft(s) with unspecified angina pectoris
I25.720	Atherosclerosis of autologous artery coronary artery bypass graft(s) with unstable angina pectoris
I25.721	Atherosclerosis of autologous artery coronary artery bypass graft(s) with angina pectoris with documented spasm
I25.728	Atherosclerosis of autologous artery coronary artery bypass graft(s) with other forms of angina pectoris
I25.729	Atherosclerosis of autologous artery coronary artery bypass graft(s) with unspecified angina pectoris
I25.730	Atherosclerosis of nonautologous biological coronary artery bypass graft(s) with unstable angina pectoris
I25.731	Atherosclerosis of nonautologous biological coronary artery bypass graft(s) with angina pectoris with documented spasm
I25.738	Atherosclerosis of nonautologous biological coronary artery bypass graft(s) with other forms of angina pectoris
I25.739	Atherosclerosis of nonautologous biological coronary artery bypass graft(s) with unspecified angina pectoris
I25.750	Atherosclerosis of native coronary artery of transplanted heart with unstable angina
I25.751	Atherosclerosis of native coronary artery of transplanted heart with angina pectoris with documented spasm
I25.758	Atherosclerosis of native coronary artery of transplanted heart with other forms of angina pectoris
I25.759	Atherosclerosis of native coronary artery of transplanted heart with unspecified angina pectoris
I25.760	Atherosclerosis of bypass graft of coronary artery of transplanted heart with unstable angina

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I25.761	Atherosclerosis of bypass graft of coronary artery of transplanted heart with angina pectoris with documented spasm
I25.768	Atherosclerosis of bypass graft of coronary artery of transplanted heart with other forms of angina pectoris
I25.769	Atherosclerosis of bypass graft of coronary artery of transplanted heart with unspecified angina pectoris
I25.790	Atherosclerosis of other coronary artery bypass graft(s) with unstable angina pectoris
I25.791	Atherosclerosis of other coronary artery bypass graft(s) with angina pectoris with documented spasm
I25.798	Atherosclerosis of other coronary artery bypass graft(s) with other forms of angina pectoris
I25.799	Atherosclerosis of other coronary artery bypass graft(s) with unspecified angina pectoris
I25.810	Atherosclerosis of coronary artery bypass graft(s) without angina pectoris
I25.811	Atherosclerosis of native coronary artery of transplanted heart without angina pectoris
I25.812	Atherosclerosis of bypass graft of coronary artery of transplanted heart without angina pectoris
Ischemic heart disease (other)	
I20.8	Other forms of angina pectoris
I20.9	Angina pectoris, unspecified
I23.7	Postinfarction angina
I24.0	Acute coronary thrombosis not resulting in myocardial infarction
I24.8	Other forms of acute ischemic heart disease
I25.2	Old myocardial infarction
I25.41	Coronary artery aneurysm
I25.42	Coronary artery dissection
I25.5	Ischemic cardiomyopathy
I25.6	Silent myocardial ischemia
I25.82	Chronic total occlusion of coronary artery
I25.83	Coronary atherosclerosis due to lipid rich plaque
I25.84	Coronary atherosclerosis due to calcified coronary lesion
I25.89	Other forms of chronic ischemic heart disease
I25.9	Chronic ischemic heart disease, unspecified

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
ST elevation (STEMI) and non-ST elevation (NSTEMI) myocardial infarction	
I21.01	ST elevation (STEMI) myocardial infarction involving left main coronary artery
I21.02	ST elevation (STEMI) myocardial infarction involving left anterior descending coronary artery
I21.09	ST elevation (STEMI) myocardial infarction involving other coronary artery of anterior wall
I21.11	ST elevation (STEMI) myocardial infarction involving right coronary artery
I21.19	ST elevation (STEMI) myocardial infarction involving other coronary artery of inferior wall
I21.21	ST elevation (STEMI) myocardial infarction involving left circumflex coronary artery
I21.29	ST elevation (STEMI) myocardial infarction involving other sites
I21.3	ST elevation (STEMI) myocardial infarction of unspecified site
I21.4	Non-ST elevation (NSTEMI) myocardial infarction
I22.0	Subsequent ST elevation (STEMI) myocardial infarction of anterior wall
I22.1	Subsequent ST elevation (STEMI) myocardial infarction of inferior wall
I22.2	Subsequent non-ST elevation (NSTEMI) myocardial infarction
I22.8	Subsequent ST elevation (STEMI) myocardial infarction of other sites
I22.9	Subsequent ST elevation (STEMI) myocardial infarction of unspecified site
Presence of cardiac and vascular implants and grafts and other postprocedural states	
Z95.1	Presence of aortocoronary bypass graft
Z95.5	Presence of coronary angioplasty implant and graft
Z98.61	Coronary angioplasty status
Occlusion and stenosis of precerebral and cerebral arteries, not resulting in cerebral infarction	
I65.01	Occlusion and stenosis of right vertebral artery
I65.02	Occlusion and stenosis of left vertebral artery
I65.03	Occlusion and stenosis of bilateral vertebral arteries
I65.09	Occlusion and stenosis of unspecified vertebral artery
I65.1	Occlusion and stenosis of basilar artery
I65.21	Occlusion and stenosis of right carotid artery
I65.22	Occlusion and stenosis of left carotid artery

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I65.23	Occlusion and stenosis of bilateral carotid arteries
I65.29	Occlusion and stenosis of unspecified carotid artery
I65.8	Occlusion and stenosis of other precerebral arteries
I65.9	Occlusion and stenosis of unspecified precerebral artery
I66.01	Occlusion and stenosis of right middle cerebral artery
I66.02	Occlusion and stenosis of left middle cerebral artery
I66.03	Occlusion and stenosis of bilateral middle cerebral arteries
I66.09	Occlusion and stenosis of unspecified middle cerebral artery
I66.11	Occlusion and stenosis of right anterior cerebral artery
I66.12	Occlusion and stenosis of left anterior cerebral artery
I66.13	Occlusion and stenosis of bilateral anterior cerebral arteries
I66.19	Occlusion and stenosis of unspecified anterior cerebral artery
I66.21	Occlusion and stenosis of right posterior cerebral artery
I66.22	Occlusion and stenosis of left posterior cerebral artery
I66.23	Occlusion and stenosis of bilateral posterior cerebral arteries
I66.29	Occlusion and stenosis of unspecified posterior cerebral artery
I66.3	Occlusion and stenosis of cerebellar arteries
I66.8	Occlusion and stenosis of other cerebral arteries
I66.9	Occlusion and stenosis of unspecified cerebral artery
Cerebrovascular diseases (other)	
I67.2	Cerebral atherosclerosis
I67.81	Acute cerebrovascular insufficiency
I67.82	Cerebral ischemia
I67.89	Other cerebrovascular disease
I67.9	Cerebrovascular disease, unspecified
I68.8	Other cerebrovascular disorders in diseases classified elsewhere

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
Cerebral infarction	
I63.00	Cerebral infarction due to thrombosis of unspecified precerebral artery
I63.011	Cerebral infarction due to thrombosis of right vertebral artery
I63.012	Cerebral infarction due to thrombosis of left vertebral artery
I63.013	Cerebral infarction due to thrombosis of bilateral vertebral arteries
I63.019	Cerebral infarction due to thrombosis of unspecified vertebral artery
I63.02	Cerebral infarction due to thrombosis of basilar artery
I63.031	Cerebral infarction due to thrombosis of right carotid artery
I63.032	Cerebral infarction due to thrombosis of left carotid artery
I63.033	Cerebral infarction due to thrombosis of bilateral carotid arteries
I63.039	Cerebral infarction due to thrombosis of unspecified carotid artery
I63.09	Cerebral infarction due to thrombosis of other precerebral artery
I63.20	Cerebral infarction due to unspecified occlusion or stenosis of unspecified precerebral arteries
I63.211	Cerebral infarction due to unspecified occlusion or stenosis of right vertebral artery
I63.212	Cerebral infarction due to unspecified occlusion or stenosis of left vertebral artery
I63.213	Cerebral infarction due to unspecified occlusion or stenosis of bilateral vertebral arteries
I63.219	Cerebral infarction due to unspecified occlusion or stenosis of unspecified vertebral artery
I63.22	Cerebral infarction due to unspecified occlusion or stenosis of basilar arteries
I63.231	Cerebral infarction due to unspecified occlusion or stenosis of right carotid arteries
I63.232	Cerebral infarction due to unspecified occlusion or stenosis of left carotid arteries
I63.239	Cerebral infarction due to unspecified occlusion or stenosis of unspecified carotid arteries
I63.29	Cerebral infarction due to unspecified occlusion or stenosis of other precerebral artery
I63.30	Cerebral infarction due to thrombosis of unspecified cerebral artery
I63.311	Cerebral infarction due to thrombosis of right middle cerebral artery
I63.312	Cerebral infarction due to thrombosis of left middle cerebral artery
I63.313	Cerebral infarction due to thrombosis of bilateral middle cerebral arteries

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I63.319	Cerebral infarction due to thrombosis of unspecified middle cerebral artery
I63.321	Cerebral infarction due to thrombosis of right anterior cerebral artery
I63.322	Cerebral infarction due to thrombosis of left anterior cerebral artery
I63.323	Cerebral infarction due to thrombosis of bilateral anterior cerebral arteries
I63.329	Cerebral infarction due to thrombosis of unspecified anterior cerebral artery
I63.331	Cerebral infarction due to thrombosis of right posterior cerebral artery
I63.332	Cerebral infarction due to thrombosis of left posterior cerebral artery
I63.333	Cerebral infarction due to thrombosis of bilateral posterior cerebral arteries
I63.339	Cerebral infarction due to thrombosis of unspecified posterior cerebral artery
I63.341	Cerebral infarction due to thrombosis of right cerebellar artery
I63.342	Cerebral infarction due to thrombosis of left cerebellar artery
I63.343	Cerebral infarction due to thrombosis of bilateral cerebellar arteries
I63.349	Cerebral infarction due to thrombosis of unspecified cerebellar artery
I63.39	Cerebral infarction due to thrombosis of other cerebral artery
I63.50	Cerebral infarction due to unspecified occlusion or stenosis of unspecified cerebral artery
I63.511	Cerebral infarction due to unspecified occlusion or stenosis of right middle cerebral artery
I63.512	Cerebral infarction due to unspecified occlusion or stenosis of left middle cerebral artery
I63.513	Cerebral infarction due to unspecified occlusion or stenosis of bilateral middle cerebral arteries
I63.519	Cerebral infarction due to unspecified occlusion or stenosis of unspecified middle cerebral artery
I63.521	Cerebral infarction due to unspecified occlusion or stenosis of right anterior cerebral artery
I63.522	Cerebral infarction due to unspecified occlusion or stenosis of left anterior cerebral artery
I63.523	Cerebral infarction due to unspecified occlusion or stenosis of bilateral anterior cerebral arteries
I63.529	Cerebral infarction due to unspecified occlusion or stenosis of unspecified anterior cerebral artery

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I63.531	Cerebral infarction due to unspecified occlusion or stenosis of right posterior cerebral artery
I63.532	Cerebral infarction due to unspecified occlusion or stenosis of left posterior cerebral artery
I63.533	Cerebral infarction due to unspecified occlusion or stenosis of bilateral posterior cerebral arteries
I63.539	Cerebral infarction due to unspecified occlusion or stenosis of unspecified posterior cerebral artery
I63.541	Cerebral infarction due to unspecified occlusion or stenosis of right cerebellar artery
I63.542	Cerebral infarction due to unspecified occlusion or stenosis of left cerebellar artery
I63.543	Cerebral infarction due to unspecified occlusion or stenosis of bilateral cerebellar arteries
I63.549	Cerebral infarction due to unspecified occlusion or stenosis of unspecified cerebellar artery
I63.59	Cerebral infarction due to unspecified occlusion or stenosis of other cerebral artery
I63.8	Other cerebral infarction
I63.9	Cerebral infarction, unspecified

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
Transient cerebral ischemic attack	
G45.8	Other transient cerebral ischemic attacks and related syndromes
G45.9	Transient cerebral ischemic attack, unspecified
Z86.73	Personal history of transient ischemic attack (TIA), and cerebral infarction without residual deficits
Vascular syndromes of brain in cerebrovascular diseases	
G46.0	Middle cerebral artery syndrome
G46.1	Anterior cerebral artery syndrome
G46.2	Posterior cerebral artery syndrome
G46.3	Brain stem stroke syndrome
G46.4	Cerebellar stroke syndrome
G46.5	Pure motor lacunar syndrome
G46.6	Pure sensory lacunar syndrome
G46.7	Other lacunar syndromes
G46.8	Other vascular syndromes of brain in cerebrovascular diseases
Atherosclerosis	
I70.0	Atherosclerosis of aorta
I70.1	Atherosclerosis of renal artery
I70.201	Unspecified atherosclerosis of native arteries of extremities, right leg
I70.202	Unspecified atherosclerosis of native arteries of extremities, left leg
I70.203	Unspecified atherosclerosis of native arteries of extremities, bilateral legs
I70.208	Unspecified atherosclerosis of native arteries of extremities, other extremity
I70.209	Unspecified atherosclerosis of native arteries of extremities, unspecified extremity
I70.211	Atherosclerosis of native arteries of extremities with intermittent claudication, right leg
I70.212	Atherosclerosis of native arteries of extremities with intermittent claudication, left leg

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I70.213	Atherosclerosis of native arteries of extremities with intermittent claudication, bilateral legs
I70.218	Atherosclerosis of native arteries of extremities with intermittent claudication, other extremity
I70.219	Atherosclerosis of native arteries of extremities with intermittent claudication, unspecified extremity
I70.221	Atherosclerosis of native arteries of extremities with rest pain, right leg
I70.222	Atherosclerosis of native arteries of extremities with rest pain, left leg
I70.223	Atherosclerosis of native arteries of extremities with rest pain, bilateral legs
I70.228	Atherosclerosis of native arteries of extremities with rest pain, other extremity
I70.229	Atherosclerosis of native arteries of extremities with rest pain, unspecified extremity
I70.231	Atherosclerosis of native arteries of right leg with ulceration of thigh
I70.232	Atherosclerosis of native arteries of right leg with ulceration of calf
I70.233	Atherosclerosis of native arteries of right leg with ulceration of ankle
I70.234	Atherosclerosis of native arteries of right leg with ulceration of heel and midfoot
I70.235	Atherosclerosis of native arteries of right leg with ulceration of other part of foot
I70.238	Atherosclerosis of native arteries of right leg with ulceration of other part of lower leg
I70.239	Atherosclerosis of native arteries of right leg with ulceration of unspecified site
I70.241	Atherosclerosis of native arteries of left leg with ulceration of thigh
I70.242	Atherosclerosis of native arteries of left leg with ulceration of calf
I70.243	Atherosclerosis of native arteries of left leg with ulceration of ankle
I70.244	Atherosclerosis of native arteries of left leg with ulceration of heel and midfoot
I70.245	Atherosclerosis of native arteries of left leg with ulceration of other part of foot
I70.248	Atherosclerosis of native arteries of left leg with ulceration of other part of lower leg
I70.249	Atherosclerosis of native arteries of left leg with ulceration of unspecified site
I70.25	Atherosclerosis of native arteries of other extremities with ulceration
I70.261	Atherosclerosis of native arteries of extremities with gangrene, right leg
I70.262	Atherosclerosis of native arteries of extremities with gangrene, left leg

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I70.263	Atherosclerosis of native arteries of extremities with gangrene, bilateral legs
I70.268	Atherosclerosis of native arteries of extremities with gangrene, other extremity
I70.269	Atherosclerosis of native arteries of extremities with gangrene, unspecified extremity
I70.291	Other atherosclerosis of native arteries of extremities, right leg
I70.292	Other atherosclerosis of native arteries of extremities, left leg
I70.293	Other atherosclerosis of native arteries of extremities, bilateral legs
I70.298	Other atherosclerosis of native arteries of extremities, other extremity
I70.299	Other atherosclerosis of native arteries of extremities, unspecified extremity
I70.301	Unspecified atherosclerosis of unspecified type of bypass graft(s) of the extremities, right leg
I70.302	Unspecified atherosclerosis of unspecified type of bypass graft(s) of the extremities, left leg
I70.303	Unspecified atherosclerosis of unspecified type of bypass graft(s) of the extremities, bilateral legs
I70.308	Unspecified atherosclerosis of unspecified type of bypass graft(s) of the extremities, other extremity
I70.309	Unspecified atherosclerosis of unspecified type of bypass graft(s) of the extremities, unspecified extremity
I70.311	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with intermittent claudication, right leg
I70.312	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with intermittent claudication, left leg
I70.313	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with intermittent claudication, bilateral legs
I70.318	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with intermittent claudication, other extremity
I70.319	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with intermittent claudication, unspecified extremity
I70.321	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with rest pain, right leg
I70.322	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with rest pain, left leg
I70.323	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with rest pain, bilateral legs
I70.328	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with rest pain, other extremity
I70.329	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with rest pain, unspecified extremity

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
170.331	Atherosclerosis of unspecified type of bypass graft(s) of the right leg with ulceration of thigh
170.332	Atherosclerosis of unspecified type of bypass graft(s) of the right leg with ulceration of calf
170.333	Atherosclerosis of unspecified type of bypass graft(s) of the right leg with ulceration of ankle
170.334	Atherosclerosis of unspecified type of bypass graft(s) of the right leg with ulceration of heel and midfoot
170.335	Atherosclerosis of unspecified type of bypass graft(s) of the right leg with ulceration of other part of foot
170.338	Atherosclerosis of unspecified type of bypass graft(s) of the right leg with ulceration of other part of lower leg
170.339	Atherosclerosis of unspecified type of bypass graft(s) of the right leg with ulceration of unspecified site
170.341	Atherosclerosis of unspecified type of bypass graft(s) of the left leg with ulceration of thigh
170.342	Atherosclerosis of unspecified type of bypass graft(s) of the left leg with ulceration of calf
170.343	Atherosclerosis of unspecified type of bypass graft(s) of the left leg with ulceration of ankle
170.344	Atherosclerosis of unspecified type of bypass graft(s) of the left leg with ulceration of heel and midfoot
170.345	Atherosclerosis of unspecified type of bypass graft(s) of the left leg with ulceration of other part of foot
170.348	Atherosclerosis of unspecified type of bypass graft(s) of the left leg with ulceration of other part of lower leg
170.349	Atherosclerosis of unspecified type of bypass graft(s) of the left leg with ulceration of unspecified site
170.35	Atherosclerosis of unspecified type of bypass graft(s) of other extremity with ulceration
170.361	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with gangrene, right leg
170.362	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with gangrene, left leg
170.363	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with gangrene, bilateral legs
170.368	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with gangrene, other extremity
170.369	Atherosclerosis of unspecified type of bypass graft(s) of the extremities with gangrene, unspecified extremity
170.391	Other atherosclerosis of unspecified type of bypass graft(s) of the extremities, right leg
170.392	Other atherosclerosis of unspecified type of bypass graft(s) of the extremities, left leg
170.393	Other atherosclerosis of unspecified type of bypass graft(s) of the extremities, bilateral legs

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
170.398	Other atherosclerosis of unspecified type of bypass graft(s) of the extremities, other extremity
170.399	Other atherosclerosis of unspecified type of bypass graft(s) of the extremities, unspecified extremity
170.401	Unspecified atherosclerosis of autologous vein bypass graft(s) of the extremities, right leg
170.402	Unspecified atherosclerosis of autologous vein bypass graft(s) of the extremities, left leg
170.403	Unspecified atherosclerosis of autologous vein bypass graft(s) of the extremities, bilateral legs
170.408	Unspecified atherosclerosis of autologous vein bypass graft(s) of the extremities, other extremity
170.409	Unspecified atherosclerosis of autologous vein bypass graft(s) of the extremities, unspecified extremity
170.411	Atherosclerosis of autologous vein bypass graft(s) of the extremities with intermittent claudication, right leg
170.412	Atherosclerosis of autologous vein bypass graft(s) of the extremities with intermittent claudication, left leg
170.413	Atherosclerosis of autologous vein bypass graft(s) of the extremities with intermittent claudication, bilateral legs
170.418	Atherosclerosis of autologous vein bypass graft(s) of the extremities with intermittent claudication, other extremity
170.419	Atherosclerosis of autologous vein bypass graft(s) of the extremities with intermittent claudication, unspecified extremity
170.421	Atherosclerosis of autologous vein bypass graft(s) of the extremities with rest pain, right leg
170.422	Atherosclerosis of autologous vein bypass graft(s) of the extremities with rest pain, left leg
170.423	Atherosclerosis of autologous vein bypass graft(s) of the extremities with rest pain, bilateral legs
170.428	Atherosclerosis of autologous vein bypass graft(s) of the extremities with rest pain, other extremity
170.429	Atherosclerosis of autologous vein bypass graft(s) of the extremities with rest pain, unspecified extremity
170.431	Atherosclerosis of autologous vein bypass graft(s) of the right leg with ulceration of thigh
170.432	Atherosclerosis of autologous vein bypass graft(s) of the right leg with ulceration of calf
170.433	Atherosclerosis of autologous vein bypass graft(s) of the right leg with ulceration of ankle
170.434	Atherosclerosis of autologous vein bypass graft(s) of the right leg with ulceration of heel and midfoot
170.435	Atherosclerosis of autologous vein bypass graft(s) of the right leg with ulceration of other part of foot

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I70.438	Atherosclerosis of autologous vein bypass graft(s) of the right leg with ulceration of other part of lower leg
I70.439	Atherosclerosis of autologous vein bypass graft(s) of the right leg with ulceration of unspecified site
I70.441	Atherosclerosis of autologous vein bypass graft(s) of the left leg with ulceration of thigh
I70.442	Atherosclerosis of autologous vein bypass graft(s) of the left leg with ulceration of calf
I70.443	Atherosclerosis of autologous vein bypass graft(s) of the left leg with ulceration of ankle
I70.444	Atherosclerosis of autologous vein bypass graft(s) of the left leg with ulceration of heel and midfoot
I70.445	Atherosclerosis of autologous vein bypass graft(s) of the left leg with ulceration of other part of foot
I70.448	Atherosclerosis of autologous vein bypass graft(s) of the left leg with ulceration of other part of lower leg
I70.449	Atherosclerosis of autologous vein bypass graft(s) of the left leg with ulceration of unspecified site
I70.45	Atherosclerosis of autologous vein bypass graft(s) of other extremity with ulceration
I70.461	Atherosclerosis of autologous vein bypass graft(s) of the extremities with gangrene, right leg
I70.462	Atherosclerosis of autologous vein bypass graft(s) of the extremities with gangrene, left leg
I70.463	Atherosclerosis of autologous vein bypass graft(s) of the extremities with gangrene, bilateral legs
I70.468	Atherosclerosis of autologous vein bypass graft(s) of the extremities with gangrene, other extremity
I70.469	Atherosclerosis of autologous vein bypass graft(s) of the extremities with gangrene, unspecified extremity
I70.491	Other atherosclerosis of autologous vein bypass graft(s) of the extremities, right leg
I70.492	Other atherosclerosis of autologous vein bypass graft(s) of the extremities, left leg
I70.493	Other atherosclerosis of autologous vein bypass graft(s) of the extremities, bilateral legs
I70.498	Other atherosclerosis of autologous vein bypass graft(s) of the extremities, other extremity
I70.499	Other atherosclerosis of autologous vein bypass graft(s) of the extremities, unspecified extremity
I70.501	Unspecified atherosclerosis of nonautologous biological bypass graft(s) of the extremities, right leg
I70.502	Unspecified atherosclerosis of nonautologous biological bypass graft(s) of the extremities, left leg

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
170.503	Unspecified atherosclerosis of nonautologous biological bypass graft(s) of the extremities, bilateral legs
170.508	Unspecified atherosclerosis of nonautologous biological bypass graft(s) of the extremities, other extremity
170.509	Unspecified atherosclerosis of nonautologous biological bypass graft(s) of the extremities, unspecified extremity
170.511	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with intermittent claudication, right leg
170.512	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with intermittent claudication, left leg
170.513	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with intermittent claudication, bilateral legs
170.518	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with intermittent claudication, other extremity
170.519	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with intermittent claudication, unspecified extremity
170.521	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with rest pain, right leg
170.522	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with rest pain, left leg
170.523	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with rest pain, bilateral legs
170.528	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with rest pain, other extremity
170.529	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with rest pain, unspecified extremity
170.531	Atherosclerosis of nonautologous biological bypass graft(s) of the right leg with ulceration of thigh
170.532	Atherosclerosis of nonautologous biological bypass graft(s) of the right leg with ulceration of calf
170.533	Atherosclerosis of nonautologous biological bypass graft(s) of the right leg with ulceration of ankle
170.534	Atherosclerosis of nonautologous biological bypass graft(s) of the right leg with ulceration of heel and midfoot
170.535	Atherosclerosis of nonautologous biological bypass graft(s) of the right leg with ulceration of other part of foot
170.538	Atherosclerosis of nonautologous biological bypass graft(s) of the right leg with ulceration of other part of lower leg
170.539	Atherosclerosis of nonautologous biological bypass graft(s) of the right leg with ulceration of unspecified site
170.541	Atherosclerosis of nonautologous biological bypass graft(s) of the left leg with ulceration of thigh
170.542	Atherosclerosis of nonautologous biological bypass graft(s) of the left leg with ulceration of calf

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I70.543	Atherosclerosis of nonautologous biological bypass graft(s) of the left leg with ulceration of ankle
I70.544	Atherosclerosis of nonautologous biological bypass graft(s) of the left leg with ulceration of heel and midfoot
I70.545	Atherosclerosis of nonautologous biological bypass graft(s) of the left leg with ulceration of other part of foot
I70.548	Atherosclerosis of nonautologous biological bypass graft(s) of the left leg with ulceration of other part of lower leg
I70.549	Atherosclerosis of nonautologous biological bypass graft(s) of the left leg with ulceration of unspecified site
I70.55	Atherosclerosis of nonautologous biological bypass graft(s) of other extremity with ulceration
I70.561	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with gangrene, right leg
I70.562	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with gangrene, left leg
I70.563	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with gangrene, bilateral legs
I70.568	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with gangrene, other extremity
I70.569	Atherosclerosis of nonautologous biological bypass graft(s) of the extremities with gangrene, unspecified extremity
I70.591	Other atherosclerosis of nonautologous biological bypass graft(s) of the extremities, right leg
I70.592	Other atherosclerosis of nonautologous biological bypass graft(s) of the extremities, left leg
I70.593	Other atherosclerosis of nonautologous biological bypass graft(s) of the extremities, bilateral legs
I70.598	Other atherosclerosis of nonautologous biological bypass graft(s) of the extremities, other extremity
I70.599	Other atherosclerosis of nonautologous biological bypass graft(s) of the extremities, unspecified extremity
I70.601	Unspecified atherosclerosis of nonbiological bypass graft(s) of the extremities, right leg
I70.602	Unspecified atherosclerosis of nonbiological bypass graft(s) of the extremities, left leg
I70.603	Unspecified atherosclerosis of nonbiological bypass graft(s) of the extremities, bilateral legs
I70.608	Unspecified atherosclerosis of nonbiological bypass graft(s) of the extremities, other extremity
I70.609	Unspecified atherosclerosis of nonbiological bypass graft(s) of the extremities, unspecified extremity
I70.611	Atherosclerosis of nonbiological bypass graft(s) of the extremities with intermittent claudication, right leg

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I70.612	Atherosclerosis of nonbiological bypass graft(s) of the extremities with intermittent claudication, left leg
I70.613	Atherosclerosis of nonbiological bypass graft(s) of the extremities with intermittent claudication, bilateral legs
I70.618	Atherosclerosis of nonbiological bypass graft(s) of the extremities with intermittent claudication, other extremity
I70.619	Atherosclerosis of nonbiological bypass graft(s) of the extremities with intermittent claudication, unspecified extremity
I70.621	Atherosclerosis of nonbiological bypass graft(s) of the extremities with rest pain, right leg
I70.622	Atherosclerosis of nonbiological bypass graft(s) of the extremities with rest pain, left leg
I70.623	Atherosclerosis of nonbiological bypass graft(s) of the extremities with rest pain, bilateral legs
I70.628	Atherosclerosis of nonbiological bypass graft(s) of the extremities with rest pain, other extremity
I70.629	Atherosclerosis of nonbiological bypass graft(s) of the extremities with rest pain, unspecified extremity
I70.631	Atherosclerosis of nonbiological bypass graft(s) of the right leg with ulceration of thigh
I70.632	Atherosclerosis of nonbiological bypass graft(s) of the right leg with ulceration of calf
I70.633	Atherosclerosis of nonbiological bypass graft(s) of the right leg with ulceration of ankle
I70.634	Atherosclerosis of nonbiological bypass graft(s) of the right leg with ulceration of heel and midfoot
I70.635	Atherosclerosis of nonbiological bypass graft(s) of the right leg with ulceration of other part of foot
I70.638	Atherosclerosis of nonbiological bypass graft(s) of the right leg with ulceration of other part of lower leg
I70.639	Atherosclerosis of nonbiological bypass graft(s) of the right leg with ulceration of unspecified site
I70.641	Atherosclerosis of nonbiological bypass graft(s) of the left leg with ulceration of thigh
I70.642	Atherosclerosis of nonbiological bypass graft(s) of the left leg with ulceration of calf
I70.643	Atherosclerosis of nonbiological bypass graft(s) of the left leg with ulceration of ankle
I70.644	Atherosclerosis of nonbiological bypass graft(s) of the left leg with ulceration of heel and midfoot
I70.645	Atherosclerosis of nonbiological bypass graft(s) of the left leg with ulceration of other part of foot
I70.648	Atherosclerosis of nonbiological bypass graft(s) of the left leg with ulceration of other part of lower leg

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I70.649	Atherosclerosis of nonbiological bypass graft(s) of the left leg with ulceration of unspecified site
I70.65	Atherosclerosis of nonbiological bypass graft(s) of other extremity with ulceration
I70.661	Atherosclerosis of nonbiological bypass graft(s) of the extremities with gangrene, right leg
I70.662	Atherosclerosis of nonbiological bypass graft(s) of the extremities with gangrene, left leg
I70.663	Atherosclerosis of nonbiological bypass graft(s) of the extremities with gangrene, bilateral legs
I70.668	Atherosclerosis of nonbiological bypass graft(s) of the extremities with gangrene, other extremity
I70.669	Atherosclerosis of nonbiological bypass graft(s) of the extremities with gangrene, unspecified extremity
I70.691	Other atherosclerosis of nonbiological bypass graft(s) of the extremities, right leg
I70.692	Other atherosclerosis of nonbiological bypass graft(s) of the extremities, left leg
I70.693	Other atherosclerosis of nonbiological bypass graft(s) of the extremities, bilateral legs
I70.698	Other atherosclerosis of nonbiological bypass graft(s) of the extremities, other extremity
I70.699	Other atherosclerosis of nonbiological bypass graft(s) of the extremities, unspecified extremity
I70.701	Unspecified atherosclerosis of other type of bypass graft(s) of the extremities, right leg
I70.702	Unspecified atherosclerosis of other type of bypass graft(s) of the extremities, left leg
I70.703	Unspecified atherosclerosis of other type of bypass graft(s) of the extremities, bilateral legs
I70.708	Unspecified atherosclerosis of other type of bypass graft(s) of the extremities, other extremity
I70.709	Unspecified atherosclerosis of other type of bypass graft(s) of the extremities, unspecified extremity
I70.711	Atherosclerosis of other type of bypass graft(s) of the extremities with intermittent claudication, right leg
I70.712	Atherosclerosis of other type of bypass graft(s) of the extremities with intermittent claudication, left leg
I70.713	Atherosclerosis of other type of bypass graft(s) of the extremities with intermittent claudication, bilateral legs
I70.718	Atherosclerosis of other type of bypass graft(s) of the extremities with intermittent claudication, other extremity
I70.719	Atherosclerosis of other type of bypass graft(s) of the extremities with intermittent claudication, unspecified extremity

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I70.721	Atherosclerosis of other type of bypass graft(s) of the extremities with rest pain, right leg
I70.722	Atherosclerosis of other type of bypass graft(s) of the extremities with rest pain, left leg
I70.723	Atherosclerosis of other type of bypass graft(s) of the extremities with rest pain, bilateral legs
I70.728	Atherosclerosis of other type of bypass graft(s) of the extremities with rest pain, other extremity
I70.729	Atherosclerosis of other type of bypass graft(s) of the extremities with rest pain, unspecified extremity
I70.731	Atherosclerosis of other type of bypass graft(s) of the right leg with ulceration of thigh
I70.732	Atherosclerosis of other type of bypass graft(s) of the right leg with ulceration of calf
I70.733	Atherosclerosis of other type of bypass graft(s) of the right leg with ulceration of ankle
I70.734	Atherosclerosis of other type of bypass graft(s) of the right leg with ulceration of heel and midfoot
I70.735	Atherosclerosis of other type of bypass graft(s) of the right leg with ulceration of other part of foot
I70.738	Atherosclerosis of other type of bypass graft(s) of the right leg with ulceration of other part of lower leg
I70.739	Atherosclerosis of other type of bypass graft(s) of the right leg with ulceration of unspecified site
I70.741	Atherosclerosis of other type of bypass graft(s) of the left leg with ulceration of thigh
I70.742	Atherosclerosis of other type of bypass graft(s) of the left leg with ulceration of calf
I70.743	Atherosclerosis of other type of bypass graft(s) of the left leg with ulceration of ankle
I70.744	Atherosclerosis of other type of bypass graft(s) of the left leg with ulceration of heel and midfoot
I70.745	Atherosclerosis of other type of bypass graft(s) of the left leg with ulceration of other part of foot
I70.748	Atherosclerosis of other type of bypass graft(s) of the left leg with ulceration of other part of lower leg
I70.749	Atherosclerosis of other type of bypass graft(s) of the left leg with ulceration of unspecified site
I70.75	Atherosclerosis of other type of bypass graft(s) of other extremity with ulceration
I70.761	Atherosclerosis of other type of bypass graft(s) of the extremities with gangrene, right leg

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
I70.762	Atherosclerosis of other type of bypass graft(s) of the extremities with gangrene, left leg
I70.763	Atherosclerosis of other type of bypass graft(s) of the extremities with gangrene, bilateral legs
I70.768	Atherosclerosis of other type of bypass graft(s) of the extremities with gangrene, other extremity
I70.769	Atherosclerosis of other type of bypass graft(s) of the extremities with gangrene, unspecified extremity
I70.791	Other atherosclerosis of other type of bypass graft(s) of the extremities, right leg
I70.792	Other atherosclerosis of other type of bypass graft(s) of the extremities, left leg
I70.793	Other atherosclerosis of other type of bypass graft(s) of the extremities, bilateral legs
I70.798	Other atherosclerosis of other type of bypass graft(s) of the extremities, other extremity
I70.799	Other atherosclerosis of other type of bypass graft(s) of the extremities, unspecified extremity
I70.8	Atherosclerosis of other arteries
I70.90	Unspecified atherosclerosis
I70.91	Generalized atherosclerosis
I70.92	Chronic total occlusion of artery of the extremities
Diseases of arteries, arterioles, and capillaries (other)	
I73.89	Other specified peripheral vascular diseases
I73.9	Peripheral vascular disease, unspecified
Arterial embolism and thrombosis	
I74.09	Other arterial embolism and thrombosis of abdominal aorta
I74.1	Embolism and thrombosis of other and unspecified parts of aorta
I74.10	Embolism and thrombosis of unspecified parts of aorta
I74.11	Embolism and thrombosis of thoracic aorta
I74.19	Embolism and thrombosis of other parts of aorta
I74.2	Embolism and thrombosis of arteries of the upper extremities
I74.3	Embolism and thrombosis of arteries of the lower extremities
I74.4	Embolism and thrombosis of arteries of extremities, unspecified

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.

 **LEQVIO**[®]
(inclisiran) injection
284 mg/1.5 mL

ICD-10-CM diagnosis code	Description
Atheroembolism	
I75.011	Atheroembolism of right upper extremity
I75.012	Atheroembolism of left upper extremity
I75.013	Atheroembolism of bilateral upper extremities
I75.019	Atheroembolism of unspecified upper extremity
I75.02	Atheroembolism of lower extremity
I75.021	Atheroembolism of right lower extremity
I75.022	Atheroembolism of left lower extremity
I75.023	Atheroembolism of bilateral lower extremities
I75.029	Atheroembolism of unspecified lower extremity
Peripheral vascular angioplasty	
Z95.820	Peripheral vascular angioplasty status with implants and grafts
Z98.62	Peripheral vascular angioplasty status
Disorders of sphingolipid metabolism and other lipid storage disorders	
E75.5	Other lipid storage disorders

Increased risk of ASCVD

Clinical risk factors	
E11	Type 2 diabetes mellitus
N18	Chronic kidney disease
I10	Essential (primary) hypertension
<p>Please note: The 3-digit ICD-10-CM codes above are not complete codes on their own. To appropriately code these diagnoses in the required format, please reference the International Classification of Diseases, Tenth Revision code manual.</p>	
Family history	
Z83.42	Family history of familial hypercholesterolemia

Reference: Centers for Medicare and Medicaid Services. ICD-10-CM. Accessed March 2, 2026. <https://www.cms.gov/medicare/coding-billing/icd-10-codes>

Please see Important Safety Information on page 32.
Click [here](#) for LEQVIO full Prescribing Information.



Indication & Important Safety Information

INDICATION

LEQVIO® (inclisiran) injection is indicated as an adjunct to diet and exercise to reduce low-density lipoprotein cholesterol (LDL-C) in adults with hypercholesterolemia, adults and pediatric patients aged 12 years and older with heterozygous familial hypercholesterolemia (HeFH), and pediatric patients aged 12 years and older with homozygous familial hypercholesterolemia (HoFH).

IMPORTANT SAFETY INFORMATION

Contraindication: LEQVIO is contraindicated in patients with a prior serious hypersensitivity reaction to inclisiran or any of the excipients in LEQVIO. Serious hypersensitivity reactions have included anaphylaxis and angioedema.

Hypersensitivity Reactions: Hypersensitivity reactions, including anaphylaxis and angioedema, have been reported in patients treated with LEQVIO. Advise patients on the signs and symptoms of hypersensitivity reactions and instruct patients to seek medical attention promptly.

Adverse Reactions: Adverse reactions in clinical trials ($\geq 3\%$ of adult patients treated with LEQVIO and more frequently than placebo) were injection site reaction, arthralgia, and bronchitis. The safety profile in pediatric patients aged 12 years and older with HeFH or HoFH was consistent with adult patients with hypercholesterolemia, except for a higher frequency of headache in pediatric patients with HeFH.

Please click [here](#) for LEQVIO full Prescribing Information.

Licensed from Alnylam Pharmaceuticals, Inc.

Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936-1080

© 2026 Novartis

3/26

FA-11607044