

For adults with moderate to severe plaque psoriasis (PsO), psoriatic arthritis (PsA), non-radiographic axial spondyloarthritis (nr-axSpA), or ankylosing spondylitis (AS).

A photograph of a man and a woman, Gary and Cyndi, dancing in a room with a white brick wall and a mirror. Gary is on the left, wearing a grey polo shirt and blue jeans, smiling and looking at Cyndi. Cyndi is on the right, wearing a colorful patterned jacket over a black top, black leggings, and black boots, also smiling and looking at Gary. The text 'MOVE, LOOK, AND FEEL BETTER.' is overlaid in large, white, bold, sans-serif capital letters across the middle of the image.

**MOVE, LOOK,
AND FEEL BETTER.**

**REAL
PATIENTS**

GARY & CYNDI

Gary is an actual patient with PsO and PsA. Cyndi is an actual patient with PsO.

This brochure contains photos of **real patients** who have taken COSENTYX® and have been compensated for their time. Individual results may vary.

INDICATIONS

COSENTYX® (secukinumab) is a prescription medicine used to treat:

- adults and children 6 years of age and older with moderate to severe plaque psoriasis (PsO) that involves large areas or many areas of the body, and who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet or UV light alone or with systemic therapy)
- adults and children 2 years of age and older with active psoriatic arthritis (PsA)
- adults and children 12 years of age and older with active ankylosing spondylitis (AS)
- adults with active non-radiographic axial spondyloarthritis (nr-axSpA) and objective signs of inflammation

IMPORTANT SAFETY INFORMATION

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

Please see Important Safety Information throughout the brochure.
Please see full Prescribing Information, including Medication Guide.

*“My elbows don’t hurt,
my joints don’t bother
me. I’m so grateful
that I found relief.”*

–Gary



REAL PsO & PsA
PATIENT

GARY

Table of Contents

About COSENTYX®	4
Psoriatic arthritis	6
Results for psoriatic arthritis	8
Psoriasis and psoriatic arthritis	10
Plaque psoriasis	12
nr-axSpA and AS	16
Results for nr-axSpA and AS	18
How COSENTYX works	22
Dosing and devices	24
Savings and support	26

IMPORTANT SAFETY INFORMATION (continued)

What is the most important information I should know about COSENTYX?

COSENTYX is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

Infections

COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have had serious infections while taking COSENTYX, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses. Some people have died from these infections.

- Your doctor should check you for TB before starting treatment with COSENTYX.
- If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with COSENTYX and during treatment with COSENTYX.
- Your doctor should watch you closely for signs and symptoms of TB during treatment with COSENTYX. **Do not use COSENTYX if you have an active TB infection.**

EXPERIENCE *matters*

COSENTYX[®] is a biologic treatment proven to help treat the symptoms of plaque psoriasis (PsO), psoriatic arthritis (PsA), non-radiographic axial spondyloarthritis (nr-axSpA), and ankylosing spondylitis (AS).

#1

COSENTYX is the #1 prescribed treatment of its kind.*

6M+

Over 6 million prescriptions have been filled in the United States.*

**18
YEARS**

Studied for over 18 years in 90 clinical trials.*

**10+
YEARS**

10 years of FDA approval.†

*Across all indications combined. As of October 2025.

†Plaque psoriasis–Jan. 2015. Psoriatic arthritis–Jan. 2016.

IMPORTANT SAFETY INFORMATION (continued)

Before starting COSENTYX, tell your doctor if you:

- are being treated for an infection
- have an infection that does not go away or that keeps coming back
- have TB or have been in close contact with someone with TB
- have or have had Hepatitis B
- think you have an infection or have symptoms of an infection such as: fevers, sweats, or chills; muscle aches; cough; shortness of breath; blood in your phlegm; weight loss; warm, red, or painful skin or sores on your body; diarrhea or stomach pain; burning when you urinate or urinate more often than normal

Please see Important Safety Information throughout the brochure.
Please see full Prescribing Information, including Medication Guide. 4



*"Five years later,
I still move and feel better."^{†§}*

REAL AS
PATIENT

MARK

[†]In a clinical trial, up to 60% of patients saw at least a 20% improvement in psoriatic arthritis symptoms at 16 weeks versus 18% of patients taking placebo.

[§]A similar percentage of people saw an improvement at 5 years as well.

 **Cosentyx[®]**
(secukinumab)

Could you be at **RISK** for permanent joint damage?

Psoriatic arthritis is an inflammatory, autoimmune condition that can affect both the joints and skin. It isn't caused from overuse of joints or aging. It's the result of an overactive immune system. Symptoms can include painful and tender joints, swollen fingers and toes, and sometimes even scaly, itchy skin. **If left untreated, psoriatic arthritis can cause permanent joint damage.**

COSENTYX® helps stop further joint damage

In a clinical trial of patients with psoriatic arthritis, about 60% of patients taking COSENTYX 150 mg or 300 mg saw at least a 20% improvement in psoriatic arthritis symptoms at 16 weeks.*

Relief that can last

84%

of people with psoriatic arthritis taking COSENTYX experienced no further joint damage at the 2-year mark.^{†‡}

*Vs 18% of patients taking placebo.

†Everyone's experience is different. These results were from a less rigorous analysis, so it is not known if symptom improvement was due to COSENTYX.

‡About 76% of patients taking COSENTYX 150 mg or 300 mg had no further joint damage at the hands, wrists, and feet at Week 24, compared to 68% taking placebo. At 2 years, 84% of patients taking COSENTYX 300 mg had no further joint damage (71% of patients taking COSENTYX 150 mg).

Please see Important Safety Information throughout the brochure. Please see full Prescribing Information, including Medication Guide. 6



"Real relief means movement, to be a dad, to be a grandpa, just to do life."

REAL PsO & PsA
PATIENT

JORGE

IMPORTANT SAFETY INFORMATION (continued)

After starting COSENTYX, call your doctor right away if you have any signs of infection listed on page 4. Do not use COSENTYX if you have any signs of infection unless you are instructed to by your doctor.

What are the possible side effects of COSENTYX?

COSENTYX may cause serious side effects, including:

Serious allergic reactions

Serious allergic reactions can occur. Get emergency medical help right away if you get any of the following symptoms: feeling faint; swelling of your face, eyelids, lips, mouth, tongue, or throat; trouble breathing or throat tightness; chest tightness; skin rash or hives (red, itchy bumps).


If you have a severe allergic reaction, do not give another injection of COSENTYX.

 **Cosentyx**[®]
(secukinumab)

COSENTYX[®]

works fast for results that can last.

COSENTYX treats the multiple symptoms of psoriatic arthritis.



*In a clinical trial of adults taking 300 mg at 16 weeks.

†In a clinical trial of adults taking COSENTYX 300 mg as their first biologic at 12 weeks.

Please see Important Safety Information throughout the brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#). 8

Watch Elizabeth's Story



REAL PsO & PsA
PATIENT

ELIZABETH

Results for
psoriatic arthritis

IMPORTANT SAFETY INFORMATION (continued)

Inflammatory bowel disease

New cases of inflammatory bowel disease or "flare-ups" can happen with COSENTYX and can sometimes be serious. If you have inflammatory bowel disease (ulcerative colitis or Crohn's disease), tell your doctor if you have worsening disease symptoms during treatment with COSENTYX or develop new symptoms of stomach pain or diarrhea.

 **Cosentyx[®]**
(secukinumab)

PLAQUE PSORIASIS *and* PSORIATIC ARTHRITIS *could be connected*

Joint pain from psoriatic arthritis and itchy skin plaques from psoriasis may not always occur together. However, symptoms from psoriatic arthritis and psoriasis may be part of the same psoriatic disease affecting different parts of the body.

Watch Jordan's Story

"I didn't realize that psoriatic arthritis and my psoriasis were connected until my doctor told me."

REAL PsO & PsA
PATIENT

JORDAN

understanding THE RISK

43%

43% of patients with moderate to severe plaque psoriasis also have psoriatic arthritis.

3X

People with nail psoriasis are almost 3x more likely to develop psoriatic arthritis.

4X

People with scalp psoriasis are almost 4x more likely to develop psoriatic arthritis.

COSENTYX treats both plaque psoriasis and psoriatic arthritis and helps stop further joint damage from psoriatic arthritis.

IMPORTANT SAFETY INFORMATION (continued)

Severe skin reactions that look like eczema can happen during treatment with COSENTYX from days to months after your first dose and can sometimes lead to hospitalization. Your doctor may temporarily stop treatment with COSENTYX if you develop severe skin reactions. Tell your doctor if you get any of the following signs or symptoms: redness or rash; itching; small bumps or patches; your skin is dry or feels like leather; blisters on the hands or feet that ooze or become crusty or skin peeling.

The most common side effects of COSENTYX include: cold symptoms, diarrhea, and upper respiratory tract infections.

These are not all of the possible side effects of COSENTYX. Call your doctor for medical advice about side effects.

 **Cosentyx**[®]
(secukinumab)

what is plaque psoriasis and its **SYMPTOMS?**

Psoriasis is an inflammatory autoimmune disease that often starts with symptoms like itchy, thick, red, scaly skin plaques. It's not a rash, and it's not contagious. And while it's not curable, it can be controlled with the proper treatment.

Psoriasis can look different across skin tones.



IMPORTANT SAFETY INFORMATION (continued)

Before using COSENTYX, tell your doctor if you:

- have any of the conditions or symptoms listed on page 4 for infections.
- have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- are allergic to latex. The removable needle cap on the COSENTYX Sensoready® pen and prefilled syringes (150 mg/mL, 75 mg/0.5 mL) contains latex.

Please see Important Safety Information throughout the brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#). 12

clear psoriasis fast with **RESULTS THAT CAN LAST**

Many adults taking COSENTYX® have seen clear or almost clear skin in as fast as 12 weeks, at 1 year, and still at 5 years.*

In a study of adults with plaque psoriasis:

skin clearance at 12 WEEKS

75%

Approximately 8 of 10 adults saw **75% clearer skin**.^{*†}

90%

Approximately 6 of 10 adults saw **90% clearer skin**.^{*‡}

in another study at 1 YEAR

100%

Approximately 6 out of 10 adults saw **100% clear skin** at 1 year.^{||}

in another study at 5 YEARS

LASTING RELIEF

Many who saw results at 1 year **maintained them at 5 years**.[§]

*In plaque psoriasis clinical trials, 6 out of 10 people taking COSENTYX 300 mg were clear or almost clear at 12 weeks vs ~3% taking placebo.

†vs ~5% patient taking placebo.

‡vs 1% taking placebo.

§At 3 years, patients knew they were getting COSENTYX, which may affect results.

||Everyone's experience is different. These results were from a less rigorous analysis, so it is not known if symptom improvement was due to COSENTYX.

PROVEN *to work*

even in troublesome areas,
like scalp and nails


Below are real patient photos taken during clinical trials.

SCALP



At Week 12, 53% of people on COSENTYX 300 mg saw 90% scalp clearance compared with 2% taking placebo.

NAILS



Nail psoriasis improved by 46% in people receiving COSENTYX 300 mg compared to an improvement of about 12% in people receiving placebo at Week 16.

Photos represent an average response with a dose of 300 mg. Individual results may vary.

COSENTYX treats both plaque psoriasis and psoriatic arthritis and helps stop further joint damage from psoriatic arthritis.

Please see Important Safety Information throughout the brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#). 14

Watch Nadia's Story



IMPORTANT SAFETY INFORMATION (continued)

- have recently received or are scheduled to receive an immunization (vaccine). People who take COSENTYX **should not** receive live vaccines. Children should be brought up to date with all vaccines before starting COSENTYX.
- have any other medical conditions and all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of your medicines to show your healthcare provider and pharmacist when you get a new medicine.

 **Cosentyx**[®]
(secukinumab)

what is **AXIAL SPONDYLOARTHRITIS (axSpA)?**

axSpA is a painful inflammatory disease that can take a serious toll on your day-to-day. Living with back pain and morning stiffness can be painful. axSpA includes two underdiagnosed conditions:

- **Non-radiographic axial spondyloarthritis (nr-axSpA)**
- **Ankylosing spondylitis (AS)**

In both conditions, inflammation in the spinal joints can cause chronic back pain, stiffness, fatigue, and reduced mobility. **If left untreated, this inflammation can cause permanent damage.**

IMPORTANT SAFETY INFORMATION (continued)

- are pregnant or plan to become pregnant. It is not known if COSENTYX can harm your unborn baby. You and your doctor should decide if you will use COSENTYX.
- are breastfeeding or plan to breastfeed. It is not known if COSENTYX passes into your breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.



REAL AS
PATIENT

WENDY

nr-axSpA and AS

RESULTS *fast.* RELIEF *that can last.*

Many people taking COSENTYX saw relief from symptoms in as little as 16 weeks, with similar results at 1 year.

nr-axSpA

40% Overall symptom improvement*^{††}

16 Weeks

1 Year

AS

3X Improvement in mobility^{‡§||}

2X More back pain relief^{‡¶}

16 Weeks

1 Year

5 YRS Many continued to feel relief at 5 years^{‡#}

16 Weeks

5 Years

*41% of people on COSENTYX 150 mg at Week 16 vs 28% taking placebo.

†38% of people on COSENTYX 150 mg at 1 year vs 19% taking placebo.

‡All results shown in clinical trials using subcutaneous (SC) administration (injection under the skin).

§Reported by those in a clinical trial taking COSENTYX and who completed a questionnaire that asked how difficult it was to perform physical activities.

||At 16 weeks, patients saw 35% improvement in mobility while taking COSENTYX vs 11% taking placebo.

¶At 16 weeks, patients taking COSENTYX saw 43% improvement in back pain vs 16% taking placebo.

#20% improvement in overall symptoms vs those taking placebo (61% vs 28%).

Please see Important Safety Information throughout the brochure.
Please see full Prescribing Information, including Medication Guide. 18



REAL AS
PATIENT

MARK

Results for
nr-axSpA and AS

IMPORTANT SAFETY INFORMATION

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

 **Cosentyx**[®]
(secukinumab)

MOVE *and* FEEL BETTER

When people taking COSENTYX® were given a survey by their doctors that measured how well their treatment was working,* the overall results showed improvements in some or all of the following symptoms of AS and nr-axSpA.



Total spinal pain



Morning stiffness



Fatigue



Tenderness (enthesitis)



Joint pain and swelling

*Bath Ankylosing Spondylitis Disease Activity Index.

Most people who took COSENTYX not only had less back pain and morning stiffness, they also saw improvement in their ability to move, and felt better overall.

IMPORTANT SAFETY INFORMATION (continued)

What is the most important information I should know about COSENTYX?

COSENTYX is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

Infections

COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have had serious infections while taking COSENTYX, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses. Some people have died from these infections.

Please see Important Safety Information throughout the brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#). 20

Watch Cat's Story

*"Before COSENTYX,
I couldn't do an essential stretch."*

REAL AS
PATIENT

CAT


- Your doctor should check you for TB before starting treatment with COSENTYX.
- If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with COSENTYX and during treatment with COSENTYX.
- Your doctor should watch you closely for signs and symptoms of TB during treatment with COSENTYX. **Do not use COSENTYX if you have an active TB infection.**

 **Cosentyx**
(secukinumab)

how does COSENTYX® work?

COSENTYX works differently by targeting and blocking IL-17A, a molecule in the body believed to play a role in inflammation in people with psoriatic arthritis, nr-axSpA, and AS.*

*The relationship between how COSENTYX works in the body and how it affects your symptoms is unknown.



Patient portrayal of how COSENTYX works.

Select your indication to see
how COSENTYX works

Explore PsA

Explore nr-axSpA

Explore AS

Explore PsO

"To see those fast results was huge for me. It's like night and day. I don't have people staring, I don't have to make excuses. Not hiding anymore is big."



REAL PsO & PsA
PATIENT

DARREN

IMPORTANT SAFETY INFORMATION (continued)

Before starting COSENTYX, tell your doctor if you:

- are being treated for an infection
- have an infection that does not go away or that keeps coming back
- have TB or have been in close contact with someone with TB

convenient DOSING *and* DEVICES

- The recommended dose of COSENTYX® for psoriatic arthritis, nr-axSpA, and AS is 150 mg*
- If you continue to have psoriatic arthritis or AS symptoms, your **rheumatologist may prescribe 300 mg** (taken as two 150-mg injections) at no additional cost for PsA and AS only
- The recommended dose of COSENTYX® for plaque psoriasis is 300 mg*

* Each 300-mg dose is given as 2 injections of 150 mg under the skin. For some people, a dose of 150 mg may be acceptable.

Getting started on COSENTYX

Once weekly for 5 weeks.



Staying on COSENTYX

Once every 4 weeks.



IMPORTANT SAFETY INFORMATION (continued)

- have or have had Hepatitis B
- think you have an infection or have symptoms of an infection such as: fevers, sweats, or chills; muscle aches; cough; shortness of breath; blood in your phlegm; weight loss; warm, red, or painful skin or sores on your body; diarrhea or stomach pain; burning when you urinate or urinate more often than normal

Please see Important Safety Information throughout the brochure. Please see full [Prescribing Information](#), including [Medication Guide](#). 24

get comfortable taking COSENTYX®

SENSOREADY® PEN



UNOREADY® PEN



NO PAIN

Most people reported no pain with the injections†

100% SATISFACTION

Up to 100% user satisfaction rating‡§

†Based on a study in which 79% of participants using the UnoReady Pen reported no pain with the injection at 12 weeks. In a study of 37 adults with active psoriatic arthritis using the Sensoready Pen, more than 90% of adults reported no pain or reaction during or after the injection.

‡Based on a study in which all 37 participants using the UnoReady Pen reported being satisfied or very satisfied with the device at 28 weeks for patients with moderate to severe plaque psoriasis.

§In a study of 37 adults with active psoriatic arthritis. Nearly 9 out of 10 adults reported being satisfied or very satisfied with self-injection using the Sensoready Pen.

Find resources on how to use the self-injection devices.

[Explore Resources](#)



IV infusion

COSENTYX also offers a monthly, 30-minute infusion for people with PsA, AS, or nr-axSpA. Plus, each infusion is administered by a healthcare provider at an infusion center or doctor's office, so you have support available to you during the process.

Learn about IV infusion, another way to take COSENTYX.

[Learn more](#)

 **Cosentyx®**
(secukinumab)

COST SHOULDN'T BE A BARRIER *between you and your treatment*

**Novartis Patient Support will always
try to help**

Novartis Patient Support is a comprehensive program designed to help you start, stay, and save on COSENTYX®.



Novartis Patient Support

- Finding answers to questions
- Navigating insurance approvals and investigations
- Directing you to injection resources so you can start with confidence



Financial Support

If you can't afford your medicine, Novartis may be able to help.



Ongoing Support

The Novartis Patient Support team can offer additional helpful resources.


\$0 Co-Pay Plus offer

The Novartis Patient Support \$0 Co-Pay Plus* Offer

If you are privately insured, you may **pay as little as \$0 on COSENTYX** with Co-Pay Plus. The Co-Pay Plus offer can help pay for out-of-pocket costs related to your prescription, including refills. See Program Terms & Conditions for eligibility, or visit www.cosentyx.com/cost. You can also reach out directly to Novartis Patient Support at 844-COSENTYX (844-267-3689) Monday-Friday, 8:00 AM-8:00 PM ET, excluding holidays.

Please see Important Safety Information throughout the brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#). 26

"Novartis Patient Support has been nothing but excellent in helping me. They make sure my insurance is set and my prescription is delivered on time."



See how Novartis Patient Support can help

***Limitations apply.** Subject to annual co-pay benefit limit. Offer not valid under Medicare, Medicaid, or any other federal or state programs. Novartis reserves the right to rescind, revoke, or amend this program without notice. Additional limitations may apply. See complete Terms & Conditions at cosentyx.com for details.



Watch Carolyn's Story

Get help with INSURANCE COVERAGE

COVERED
until you're
COVERED

If your insurance coverage is delayed, you shouldn't have to wait. Covered Until You're Covered, also known as The Bridge Program*, can provide up to 2 years of COSENTYX® for **free** while seeking coverage.

Once your insurance is approved, your Co-Pay Plus offer may continue to help with out-of-pocket costs.

IV infusion


COSENTYX IV infusions may be covered under Medicare Part B for people with PsA, AS, and nr-axSpA. Be sure to check with your insurance provider and talk to your doctor to understand your coverage and treatment options.

*The Bridge Program applies to COSENTYX Subcutaneous Injection only. Eligible patients must have private insurance and a valid prescription for COSENTYX, and a prior authorization, predetermination, or medical exception that has been denied. Program requires the submission of an appeal of the coverage denial within the first 90 days of enrollment to remain eligible. Program provides COSENTYX for free to eligible patients for up to 2 years, or until they receive insurance coverage approval, whichever occurs earlier. A valid prescription consistent with FDA approved labeling is required. Program is not available to patients whose medications are reimbursed in whole or in part by Medicare, Medicaid, TRICARE, or any other federal or state program. Patients may be asked to reverify insurance coverage status during the course of the program. No purchase necessary. Program is not health insurance, nor is participation a guarantee of insurance coverage. Additional Limitations may apply. Novartis Pharmaceuticals Corporation reserves the right to rescind, revoke, or amend this Program without notice.

REAL AS
PATIENT

CAROLYN

**MOVE, LOOK
AND FEEL
BETTER.**



**REAL PsO & PsA
PATIENT**

GARY

**Ask your rheumatologist if COSENTYX[®]
is right for you.**



Learn more at [**COSENTYX.com**](https://www.cosentyx.com)



[**@COSENTYX \(secukinumab\)**](https://www.facebook.com/COSENTYX)



[**@COSENTYX \(secukinumab\)**](https://www.instagram.com/COSENTYX)

Please see Important Safety Information throughout the brochure.
Please see full Prescribing Information, including Medication Guide.

 **NOVARTIS**

Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936-1080