

FOR ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER
WITH MODERATE TO SEVERE HIDRADENITIS SUPPURATIVA (HS)



REAL RELIEF FROM HS SYMPTOMS FOR ADULTS AND CHILDREN 12+

Patient portrayal.
Individual results may vary.

This brochure contains photos and quotes of actual adult patients who have taken COSENTYX® and have been compensated for their time. Individual results may vary.

INDICATIONS

COSENTYX® (secukinumab) is a prescription medicine used to treat:

- adults and children 6 years of age and older with moderate to severe plaque psoriasis (PsO) that involves large areas or many areas of the body, and who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet or UV light alone or with systemic therapy)
- adults and children 2 years of age and older with active psoriatic arthritis (PsA)
- adults and children 12 years of age and older with active ankylosing spondylitis (AS)
- adults with active non-radiographic axial spondyloarthritis (nr-axSpA) and objective signs of inflammation
- children 4 years of age and older with active enthesitis-related arthritis (ERA)
- adults and children 12 years of age and older with moderate to severe hidradenitis suppurativa (HS)

IMPORTANT SAFETY INFORMATION

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

Please see Important Safety Information throughout brochure.

Please see full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.



Patient portrayal.

Table of Contents

About HS	6
Results in Adults	10
About COSENTYX®	16
Safety	18
Dosing	19
Injecting	20
Visit Prep	22
Support	24

IMPORTANT SAFETY INFORMATION (cont) What is the most important information I should know about COSENTYX?

COSENTYX is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

Infections
COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have had serious infections while taking COSENTYX, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses. Some people have died from these infections.

- Your doctor should check you for TB before starting treatment with COSENTYX.
- If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with COSENTYX and during treatment with COSENTYX.
- Your doctor should watch you closely for signs and symptoms of TB during treatment with COSENTYX.
Do not use COSENTYX if you have an active TB infection.

COSENTYX[®] (secukinumab) offers real relief for adults and **CHILDREN 12+** living with moderate to severe HS

COSENTYX, which is approved to treat adults and children 12+, targets and blocks IL-17A,* an important molecule that may contribute to inflammation that leads to HS symptoms.

**The safety of COSENTYX is well studied
— with 18 years of clinical testing.[†]**

HS diagnosis can take years—and during this time, the condition can progress. Don't wait to talk to your doctor about treatment options and ask if COSENTYX is right for your child.

*The relationship between how COSENTYX works in the body and how it affects your symptoms is unknown.

[†]Across all indications combined. Does not include patients under 18 with HS.

IMPORTANT SAFETY INFORMATION (cont)

Before starting COSENTYX, tell your doctor if you:

- are being treated for an infection
- have an infection that does not go away or that keeps coming back
- have TB or have been in close contact with someone with TB
- have or have had Hepatitis B
- think you have an infection or have symptoms of an infection such as: fevers, sweats, or chills; muscle aches; cough; shortness of breath; blood in your phlegm; weight loss; warm, red, or painful skin or sores on your body; diarrhea or stomach pain; burning when you urinate or urinate more often than normal

Please see Important Safety Information throughout brochure.
Please see full Prescribing Information, including [Medication Guide](#).

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.

"When I was a teen, it was very challenging. I would tell my younger self that there is going to be something that's going to help."



REAL
PATIENT

LILY

 **Cosentyx[®]**
(secukinumab)

HIDRADENITIS SUPPURATIVA (HS) is an inflammatory skin condition that may begin around puberty

While the exact cause is unknown, it is thought that HS symptoms start with a blocked hair follicle that triggers the immune system, producing too many inflammatory molecules, like IL-17A. In people with HS, an overactive immune system and too much IL-17A are thought to contribute to excess inflammation in the body that may cause bumps and abscesses that can hurt.

IMPORTANT SAFETY INFORMATION (cont)

After starting COSENTYX, call your doctor right away if you have any signs of infection listed on page 4.

Do not use COSENTYX if you have any signs of infection unless you are instructed to by your doctor.

What are the possible side effects of COSENTYX?

COSENTYX may cause serious side effects, including:

Serious allergic reactions

Serious allergic reactions can occur. Get emergency medical help right away if you get any of the following symptoms: feeling faint; swelling of your face, eyelids, lips, mouth, tongue, or throat; trouble breathing or throat tightness; chest tightness; skin rash or hives (red, itchy bumps).

HS can make your **CHILD** push things away

Living with HS can be a long and frustrating journey. Often, dealing with painful HS symptoms can make everyday activities a challenge—even something as simple as a hug can be too painful.




Patient portrayal.

what HS can look and feel like

HS signs and symptoms can include painful bumps or abscesses, mostly in areas where skin rubs against skin. HS abscesses can drain and may lead to tunnels that can leave scars.

Where HS symptoms commonly appear



how HS may progress

Over time, HS may progress or get worse if left untreated, and may lead to scarring.

HS can be classified into 3 stages* based on the severity of the symptoms:



MILD[†] is when there is 1 bump or multiple inflammatory bumps and abscesses, with no scarring or tunnel (sinus tract) formation.



MODERATE is when there are multiple abscesses, along with limited tunnels and/or scarring.



SEVERE is when there are multiple or extensive interconnected tunnels, abscesses, and scarring.

All images used with permission from ©DermNet NZ.

*In classifying the severity of your condition, your dermatologist may call these Hurley stages.

[†]COSENTYX® is approved to treat moderate to severe HS.

IMPORTANT SAFETY INFORMATION (cont)

If you have a severe allergic reaction, do not give another injection of COSENTYX.

Inflammatory bowel disease

New cases of inflammatory bowel disease or “flare-ups” can happen with COSENTYX and can sometimes be serious. If you have inflammatory bowel disease (ulcerative colitis or Crohn’s disease), tell your doctor if you have worsening disease symptoms during treatment with COSENTYX or develop new symptoms of stomach pain or diarrhea.

Please see Important Safety Information throughout brochure.
Please see full Prescribing Information, including Medication Guide.

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.

IMPORTANT SAFETY INFORMATION (cont)

Severe skin reactions that look like eczema can happen during treatment with COSENTYX from days to months after your first dose and can sometimes lead to hospitalization. Your doctor may temporarily stop treatment with COSENTYX if you develop severe skin reactions. Tell your doctor if you get any of the following signs or symptoms: redness or rash; itching; small bumps or patches; your skin is dry or feels like leather; blisters on the hands or feet that ooze or become crusty or skin peeling.

PROVEN RELIEF for adults with HS



REDUCTION

At least a 50% reduction of bumps and abscesses.



NO INCREASE

No increase in these symptoms.

In the 2 clinical trials, 41% and 43% of adults taking COSENTYX®* had at least a 50% reduction in the number of inflammatory bumps and abscesses, with no increase in the number of abscesses and/or draining tunnels at 16 weeks vs 29% and 26% taking placebo.

* COSENTYX 300 mg every 4 weeks (after 5 initial weekly doses).

IMPORTANT SAFETY INFORMATION (cont)

The most common side effects of COSENTYX include: cold symptoms, diarrhea, and upper respiratory tract infections.

These are not all of the possible side effects of COSENTYX. Call your doctor for medical advice about side effects.

Before using COSENTYX, tell your doctor if you:

- have any of the conditions or symptoms listed on page 4 for infections.
- have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- are allergic to latex. The removable needle cap on the COSENTYX Sensoready® pen and prefilled syringes (150 mg/mL, 75 mg/0.5 mL) contains latex.
- have recently received or are scheduled to receive an immunization (vaccine). People who take COSENTYX **should not** receive live vaccines. Children should be brought up to date with all vaccines before starting COSENTYX.

Please see Important Safety Information throughout brochure.
Please see full Prescribing Information, including Medication Guide.

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.

Watch real people with HS
talk about their experiences
taking COSENTYX

Watch now

"I have tried some of everything to manage my symptoms when I was younger. COSENTYX came in and gave me the help I needed."




REAL
PATIENT

FRANK

Results in Adults

 **Cosentyx**
(secukinumab)

ZERO FLARES *for the majority of people*



In 1 of 2 clinical trials, 75% of adults taking COSENTYX® had zero flares at Week 16.

In the same clinical trial, 25% of adults taking COSENTYX* had 1 or more flares at Week 16 vs 36% taking placebo. In the second trial, the same results were not seen.

A flare was defined as a greater than 25% increase in the number of inflammatory bumps and abscesses, with a minimum increase of 2 inflammatory bumps or abscesses.

*COSENTYX 300 mg every 4 weeks (after 5 initial weekly doses).

IMPORTANT SAFETY INFORMATION (cont)

Before using COSENTYX, tell your doctor if you:

- have any other medical conditions and all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of your medicines to show your healthcare provider and pharmacist when you get a new medicine.
- are pregnant or plan to become pregnant. It is not known if COSENTYX can harm your unborn baby. You and your doctor should decide if you will use COSENTYX.
- are breastfeeding or plan to breastfeed. It is not known if COSENTYX passes into your breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

Please see Important Safety Information throughout brochure. Please see full Prescribing Information, including Medication Guide.

Visit COSENTYX.com for more information.

Watch real people with HS talk about their experiences taking COSENTYX

Watch now



"Since finding relief with COSENTYX, it's kind of allowed my skin to breathe. I don't have to worry about wearing tank tops."

REAL PATIENT

LILY

IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about COSENTYX?

COSENTYX is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

Infections

COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have had serious infections while taking COSENTYX, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses. Some people have died from these infections.

RELIEF FROM HS SYMPTOMS *in 2 weeks and years after*



IN AS EARLY AS 2 WEEKS, many adults saw a nearly 50% reduction in bumps/abscesses.

These results were from a less rigorous analysis, so it is not known if symptom improvement was due to COSENTYX®. Everyone's experience is different.



RELIEF FROM HS AT 1 YEAR*

- Adults taking COSENTYX maintained the reduction of bumps and abscesses
- Approximately, 61% of people had no increase in tunnels
- Approximately, 73% of people had no flares

In the 2 clinical trials, at 1 year, 58% and 65% had at least a 50% reduction in the number of inflammatory bumps and abscesses, with no increase in the number of abscesses and/or draining tunnels. In addition, 72% and 74% did not experience a flare. In the clinical trials, a flare was defined as a greater than 25% increase in the number of inflammatory bumps and abscesses, with a minimum increase of 2 inflammatory bumps or abscesses.

These results were from a less rigorous analysis, so it is not known if symptom improvement was due to COSENTYX. Everyone's experience is different.

*Results are from a 300-mg dose of COSENTYX every 4 weeks (after 5 initial weekly doses). Results may vary. Everyone responds to treatment differently.

Please see Important Safety Information throughout brochure.
Please see full Prescribing Information, including Medication Guide.

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.



CONTINUED RESULTS AT 2 YEARS†

Adults taking COSENTYX maintained the reduction of bumps and abscesses.

In an extension study, which followed patients from 2 clinical trials, 76.5% and 87.5% of patients who had previously experienced at least a 50% reduction in the number of inflammatory bumps and abscesses, with no increase in the number of abscesses and/or draining tunnels at Year 1 maintained this reduction at Year 2.

Because these results were from a less rigorous analysis, no conclusions can be drawn.

†Results are from a 300-mg dose of COSENTYX every 2 weeks or every 4 weeks respectively. Results may vary. Everyone responds to treatment differently.

IMPORTANT SAFETY INFORMATION (cont)

Infections (continued)

- Your doctor should check you for TB before starting treatment with COSENTYX.
- If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with COSENTYX and during treatment with COSENTYX.
- Your doctor should watch you closely for signs and symptoms of TB during treatment with COSENTYX. **Do not use COSENTYX if you have an active TB infection.**



COSENTYX® is a biologic with over 6 million prescriptions filled* in the United States

What is a biologic?

A biologic is a protein-based medication that is made from living sources and is taken by injection or infusion.

How does COSENTYX work?

COSENTYX works within the body's immune system to selectively target and block just the IL-17A molecule, which is believed to cause inflammation throughout the body when it is overproduced. IL-17A is one of the multiple molecules that is believed to play a role in causing inflammation.

The relationship between how COSENTYX works in the body and how it affects your symptoms is unknown.

Science you can trust. Relief you can feel.

COSENTYX has been studied for more than 18 years in dozens of clinical trials across all indications.*

- Well-studied safety profile since launch in 2015
- No routine lab monitoring required during treatment

*Across all indications combined. Does not include patients under 18 with HS.

Please see Important Safety Information throughout brochure.
Please see full Prescribing Information, including Medication Guide.

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.

16

COSENTYX works DIFFERENTLY



See how
COSENTYX works

Watch now

About COSENTYX®

Patient portrayal of how COSENTYX works.

See how real patients
treat and manage HS

Watch now

IMPORTANT SAFETY INFORMATION (cont)

Before starting COSENTYX, tell your doctor if you:

- are being treated for an infection
- have an infection that does not go away or that keeps coming back
- have TB or have been in close contact with someone with TB
- Have or have had Hepatitis B
- think you have an infection or have symptoms of an infection such as: fevers, sweats, or chills; muscle aches; cough; shortness of breath; blood in your phlegm; weight loss; warm, red, or painful skin or sores on your body; diarrhea or stomach pain; burning when you urinate or urinate more often than normal

 **Cosentyx®**
(secukinumab)

17

people stick with **COSENTYX**[®]

Less than 3% of adults in clinical studies stopped taking COSENTYX due to side effects.*

* In a study of patients with moderate to severe HS.



Patient portrayal.
Individual results may vary.

IMPORTANT SAFETY INFORMATION (cont)

After starting COSENTYX, call your doctor right away if you have any signs of infection listed on page 4.

Do not use COSENTYX if you have any signs of infection unless you are instructed to by your doctor.

What are the possible side effects of COSENTYX?

COSENTYX may cause serious side effects, including:

Serious allergic reactions

Serious allergic reactions can occur. Get emergency medical help right away if you get any of the following symptoms: feeling faint; swelling of your face, eyelids, lips, mouth, tongue, or throat; trouble breathing or throat tightness; chest tightness; skin rash or hives (red, itchy bumps).

Please see Important Safety Information throughout brochure.
Please see full Prescribing Information, including Medication Guide.

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.

dosing for **CHILDREN 12+** based on body weight, helping families start treatment earlier

Your child's doctor will prescribe the dose of COSENTYX that's right for them, guided by their weight. To start, they will take COSENTYX once a week for 5 weeks, then just once a month for maintenance. This can mean fewer interruptions in both you and your child's daily routine.

for those weighing

66 to less than 198 lbs:

the recommended dose is 150 mg

- If the prescribed dose of **COSENTYX is 150 mg**, you must give 1 injection of the COSENTYX 150-mg Sensoready[®] Pen or 150-mg prefilled syringe for each dose

198 lbs or more:

the recommended dose is 300 mg

- If the prescribed dose of **COSENTYX is 300 mg**, you must give 1 injection of the COSENTYX 300-mg UnoReady[®] Pen or 2 injections of the COSENTYX 150-mg Sensoready[®] Pen

Monthly dose equals 1 dose every 4 weeks.

get comfortable with **COSENTYX®**

With once-monthly dosing, COSENTYX can help make it more convenient to manage your child's HS. And there is no routine lab monitoring during treatment.

Children should not inject themselves with COSENTYX. An adult caregiver should prepare and inject COSENTYX after receiving training from a healthcare provider on the right way to prepare and inject COSENTYX.

Talk to your doctor if you have any questions.

Injection resources

Read the detailed "Instructions for Use" that comes with your COSENTYX for information on how to prepare and inject a dose of COSENTYX, and how to properly throw away (dispose of) used COSENTYX. For added support, a Novartis Patient Support Team Member is also available at 1-844-COSENTYX (1-844-267-3689).

IMPORTANT SAFETY INFORMATION (cont)

If you have a severe allergic reaction, do not give another injection of COSENTYX.

Inflammatory bowel disease

New cases of inflammatory bowel disease or "flare-ups" can happen with COSENTYX, and can sometimes be serious. If you have inflammatory bowel disease (ulcerative colitis or Crohn's disease), tell your doctor if you have worsening disease symptoms during treatment with COSENTYX or develop new symptoms of stomach pain or diarrhea.

Sensoready® Pen



UnoReady® Pen



NO PAIN

Most people reported no pain with the injections*

**100%
SATISFACTION**

Up to 100% user satisfaction rating^{††}

*Based on a study in which 79% of participants using the UnoReady Pen reported no pain with the injection at 12 weeks. In a study of 37 adults with active psoriatic arthritis using the Sensoready Pen, more than 90% of adults reported no pain or reaction during or after the injection.

[†]Based on a study in which all 37 participants using the UnoReady Pen reported being satisfied or very satisfied with the device at 28 weeks for patients with moderate to severe plaque psoriasis.

^{††}In a study of 37 adults with active psoriatic arthritis. Nearly 9 out of 10 adults reported being satisfied or very satisfied with self-injection using the Sensoready Pen.

**Watch our supplemental
injection demonstration
videos**

Watch now

IMPORTANT SAFETY INFORMATION (cont)

Severe skin reactions that look like eczema can happen during treatment with COSENTYX from days to months after your first dose and can sometimes lead to hospitalization. Your doctor may temporarily stop treatment with COSENTYX if you develop severe skin reactions. Tell your doctor if you get any of the following signs or symptoms: redness or rash; itching; small bumps or patches; your skin is dry or feels like leather; blisters on the hands or feet that ooze or become crusty or skin peeling.



unsure what to expect at your dermatologist visit? **DON'T WORRY, WE'RE HERE TO HELP**

BEFORE THE VISIT

- Prepare your kid or teen emotionally by saying something like, "Let's decide together what you want the doctor to know."
- Keep a simple log of any new symptoms, activities, or products that seem to trigger flares
- Consider bringing photos of affected skin areas
- Make a list of everything you've tried, including over-the-counter treatments, soaps or deodorants, clothing changes, or any lifestyle adjustments

DURING THE VISIT

- The dermatologist may ask about symptoms, pain, and daily impacts
- The dermatologist will gently examine affected areas
- The dermatologist will discuss ongoing monitoring and follow-up



Please see Important Safety Information throughout brochure.
Please see full Prescribing Information, including Medication Guide.

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.

here are some **STARTER QUESTIONS** to ask your dermatologist

Understanding the Diagnosis

- How serious is it or what stage of HS does my kid/teen have?
- How do you know it's HS and not something else?

Treatment Option

- What treatments are available for a kid/teen at this age?
- What results should we expect—and how long might it take to see results?

Pain & Flare Management

- What can we do at home to help during a flare?
- When should we call you?

Lifestyle Guidance

- Are there clothing, activities, or products that may help reduce irritation?
- How can we support our child?

The Long-Term Picture

- How often should we follow up?
- How can we track progress together?

what is Novartis Patient Support?

Novartis Patient Support is a comprehensive program designed to help your child start, stay, and save on COSENTYX®.

Novartis Patient Support is not a clinical service and does not replace guidance from your healthcare provider. Our goal is to help you feel informed about your treatment from day one.

For caregivers of patients who are starting or taking COSENTYX, here's what you will get once you sign up:



Novartis Patient Support

The Novartis Patient Support team can help with:

- Finding answers to questions
- Navigating insurance approvals and investigations
- Directing you to injection resources so you can start with confidence



Financial Support

If you can't afford your medicine, Novartis may be able to help.



Co-Pay Plus offer

COVERED
until you're
COVERED

Privately insured patients may be eligible to pay as little as \$0*

Up to 2 years of COSENTYX for free with the Covered Until You're Covered program† if a patient's insurance coverage is denied



Ongoing Support

The Novartis Patient Support team can offer additional helpful resources.



To sign up for Novartis Patient Support and start receiving these free resources, call us at **1-844-COSENTYX (1-844-267-3689)** or visit **[COSENTYX.com](https://www.cosentyx.com)**.

***Limitations apply.** Subject to annual co-pay benefit limit. Offer not valid under Medicare, Medicaid, or any other federal or state programs. Novartis reserves the right to rescind, revoke, or amend this program without notice. Additional limitations may apply. See complete Terms & Conditions at <https://www.cosentyx.com/all/treatment-cost#terms-and-conditions> for details.

†Covered Until You're Covered also known as Bridge Program applies to COSENTYX Subcutaneous Injection only. Eligible patients must have private insurance and a valid prescription for COSENTYX, and a prior authorization, predetermination, or medical exception that has been denied. Program requires the submission of an appeal of the coverage denial within the first 90 days of enrollment to remain eligible. Program provides COSENTYX for free to eligible patients for up to 2 years, or until they receive insurance coverage approval, whichever occurs earlier. A valid prescription consistent with FDA-approved labeling is required. Program is not available to patients whose medications are reimbursed in whole or in part by Medicare, Medicaid, TRICARE, or any other federal or state program. Patients may be asked to reverify insurance coverage status during the course of the program. No purchase necessary. Program is not health insurance, nor is participation a guarantee of insurance coverage. Additional Limitations may apply. Novartis Pharmaceuticals Corporation reserves the right to rescind, revoke, or amend this Program without notice.

getting COSENTYX® covered

A Novartis Patient Support Team Member will work with you and your healthcare provider to help get your child's COSENTYX prescription approved. Go to [COSENTYX.com/how-to-get-cosentyx](https://www.cosentyx.com/how-to-get-cosentyx) for a full walk-through of the steps for getting COSENTYX.

GETTING COSENTYX DELIVERED

COSENTYX is sent to you from a specialty pharmacy since it's a biologic medication, which means it requires special handling and must be refrigerated.

GETTING FINANCIAL SUPPORT

COSENTYX Co-Pay Plus offer*:

Pay as little as \$0 Co-Pay for eligible patients with private insurance. In fact, 84% of people who used the COSENTYX Co-Pay Plus Program in 2024 paid nothing out of pocket for their prescription.

HELP NAVIGATING INSURANCE

In the unlikely event that your child's Private Insurance is denied, we'll do everything we can to help. Learn more about our Covered Until You're Covered program.[†] The program offers up to 2 years of COSENTYX for free while their healthcare provider tries to secure coverage. For more coverage assistance, head to [COSENTYX.com/how-to-get-cosentyx](https://www.cosentyx.com/how-to-get-cosentyx).



Learn more about how you
can get COSENTYX from
your specialty pharmacy

Learn more

***Limitations apply.** Subject to annual co-pay benefit limit. Offer not valid under Medicare, Medicaid, or any other federal or state programs. Novartis reserves the right to rescind, revoke, or amend this program without notice. Additional limitations may apply. See complete Terms & Conditions at <https://www.cosentyx.com/all/treatment-cost#terms-and-conditions> for details.

[†]Covered Until You're Covered also known as Bridge Program applies to COSENTYX Subcutaneous Injection only. Eligible patients must have private insurance and a valid prescription for COSENTYX, and a prior authorization, predetermination, or medical exception that has been denied. Program requires the submission of an appeal of the coverage denial within the first 90 days of enrollment to remain eligible. Program provides COSENTYX for free to eligible patients for up to 2 years, or until they receive insurance coverage approval, whichever occurs earlier. A valid prescription consistent with FDA-approved labeling is required. Program is not available to patients whose medications are reimbursed in whole or in part by Medicare, Medicaid, TRICARE, or any other federal or state program. Patients may be asked to reverify insurance coverage status during the course of the program. No purchase necessary. Program is not health insurance, nor is participation a guarantee of insurance coverage. Additional Limitations may apply. Novartis Pharmaceuticals Corporation reserves the right to rescind, revoke, or amend this Program without notice.



ASK YOUR CHILD'S DOCTOR ABOUT COSENTYX[®] TODAY



Visit [COSENTYX.com/hidradenitis-suppurativa/index](https://www.cosentyx.com/hidradenitis-suppurativa/index)



[@COSENTYX \(secukinumab\)](https://www.instagram.com/cosentyx)



[@COSENTYX \(secukinumab\)](https://www.facebook.com/cosentyx)

Please see Important Safety Information throughout brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.

 **NOVARTIS**

Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936-1080