




# A GUIDE TO. PLUVICTO

What to expect and  
how to prepare



**PLUVICTO<sup>®</sup>**

lutetium Lu 177 vipivotide tetraxetan  
INJECTION FOR INTRAVENOUS USE

Patient portrayal.

## What is PLUVICTO<sup>®</sup> (lutetium Lu 177 vipivotide tetraxetan)?

PLUVICTO is a prescription treatment used to treat adults with metastatic prostate-specific membrane antigen-positive (PSMA+) prostate cancer and are:

- androgen pathway modulation-naïve or -sensitive (mAPMN/S) in combination with hormone therapy, or
- androgen pathway modulation-resistant (mAPMR) after hormone therapy, and
  - appropriate to delay chemotherapy, or
  - have received prior chemotherapy

## IMPORTANT SAFETY INFORMATION

### What is the most important information I should know about PLUVICTO?

**Use of PLUVICTO involves exposure to radioactivity.**

Long-term, accruing radiation exposure is associated with an increased risk for cancer. Drink plenty of water and urinate as often as possible during the first hours after administration.

**Please see additional Important Safety Information throughout.**

This guide is your companion from start to finish.

**Your health care team** [Page 3](#)

**Key steps** [Page 4](#)

**How to prepare** [Page 5](#)

**Planning for each dose** [Pages 6–10](#)

**Novartis Patient Support™** [Page 12](#)



Patient portrayal.

## IMPORTANT SAFETY INFORMATION (continued)

### What is the most important information I should know about PLUVICTO? (continued)

To minimize radiation exposure to others following administration of PLUVICTO, limit close contact (less than 3 feet) with household contacts for 2 days or with children and pregnant women for 7 days. Refrain from sexual activity for 7 days, and sleep in a separate bedroom from household contacts for 3 days, from children for 7 days, or from pregnant women for 15 days.

**Please see additional Important Safety Information throughout.**

Throughout your PLUVICTO treatment, you will have a team of health care professionals who will work together to support you.



Doctor portrayals.



### Your ongoing care doctor

This may be a medical oncologist or urologist who **manages your metastatic prostate cancer and refers you to your treating team** if PLUVICTO could be your next step.

### Your treating team

- Your doctor and team will **deliver and oversee PLUVICTO at a treatment center**
- Nurses and other health care professionals will also coordinate your care with your doctor

## IMPORTANT SAFETY INFORMATION (continued)

**PLUVICTO may cause serious side effects, including:**

**Low level of blood cell counts, which may be severe and life-threatening.** Tell your doctor right away if you develop any new or worsening symptoms, including:

- Tiredness or weakness
- Pale skin
- Shortness of breath
- Bleeding or bruising more easily than normal or difficulty stopping bleeding
- Frequent infections with signs such as fever, chills, sore throat, or mouth ulcers



# Your PLUVICTO treatment roadmap

Patient portrayal.

## Decide with your doctor to start PLUVICTO.

If you haven't received a PSMA-PET scan recently, ask your doctor if you need to get one.

## Select a treatment center with your doctor and schedule a consult and pretreatment bloodwork there.


See [page 11](#) for help finding a treatment center.

## Get your PLUVICTO doses.

PLUVICTO is given every 6 weeks for up to 6 doses.\* After that, your PLUVICTO treatment is complete. If PLUVICTO is given together with other treatments, you may continue taking them.

[Click here to watch a video about what to expect during your journey at PLUVICTO.com.](#)

## PLUVICTO (intravenous injection or infusion)



Follow your care team's instructions for when to receive your bloodwork and each PLUVICTO dose.

\*Dosing may change or stop based on disease progression or unmanageable side effects.

PET, positron emission tomography; PSMA, prostate-specific membrane antigen.

## IMPORTANT SAFETY INFORMATION (continued)

### PLUVICTO may cause serious side effects, including (continued):

**Kidney problems, which can be severe.** You should stay well hydrated before and after treatment. Tell your doctor right away if you develop any new or worsening urinary symptoms.

All radiopharmaceuticals, including PLUVICTO, have the potential to **cause harm to an unborn baby**.

- You should use effective contraception during treatment with PLUVICTO and for 14 weeks after your last dose. PLUVICTO may cause temporary or permanent **infertility**.

**Please see additional Important Safety Information throughout.**

 **PLUVICTO®**  
lutetium Lu 177 vipivotide tetraxetan  
INJECTION FOR INTRAVENOUS USE

## Prepare for your first PLUVICTO dose

### Set your expectations

Before you go, ask your treatment center about what to expect on the day of treatment and how long treatment will take. It may take less than 1 hour.

### Get ready for your treatment

Download a playlist, podcast, game, or show to bring with you on the day of treatment and dress comfortably.

### Plan your transportation

Keep in mind that you'll need to practice social distancing after treatment and for the following days.

See [page 9](#) for more information.

Patient portrayal.

## IMPORTANT SAFETY INFORMATION (continued)

### The most common side effects of PLUVICTO include:

- Decreased blood cell counts
- Laboratory abnormalities
- Tiredness
- Dry mouth
- Nausea
- Kidney problems

Please see additional Important Safety Information throughout.

## What targeted radiation with PLUVICTO means for you

PLUVICTO is a targeted therapy with the goal to deliver radiation directly into PSMA-positive (PSMA+) cells to destroy them from within.

### After treatment, radiation does not stay in your body for long



**Within 2 days**, over half the radiation will leave your body.

**Within about 14 days**, most of the radiation (>99%) will be gone.

### Radiation exposure to others is limited and can be compared to taking 3 flights across the United States



After you receive 1 dose of PLUVICTO, care partners who follow radiation safety requirements (see [page 8](#) for more on this) may be exposed to the same amount of radiation as someone flying 3 times from the East Coast to the West Coast.

PSMA, prostate-specific membrane antigen.

## IMPORTANT SAFETY INFORMATION (continued)

These are not all of the possible side effects of PLUVICTO. Call your doctor for advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.



## What you can do after treatment

It's important to know that your sweat, blood, and urine give off radiation after each dose of PLUVICTO.



You can help eliminate this radiation through frequent urination. Your doctor may recommend drinking about 8 ounces of fluid every hour on the day of your dose to help you urinate frequently.



Patient portrayal.

### IMPORTANT SAFETY INFORMATION

#### What is the most important information I should know about PLUVICTO?

##### Use of PLUVICTO involves exposure to radioactivity.

Long-term, accruing radiation exposure is associated with an increased risk for cancer. Drink plenty of water and urinate as often as possible during the first hours after administration.

**Please see additional Important Safety Information throughout.**

## Minimize radiation exposure to others

### Keep a safe distance from others after each dose



Stay more than 3 feet away, about an arm's length, from:

- Adults for 2 days
- Children, pregnant women, or pets for 7 days



Sleep alone in a separate room, away from:

- Adults and pets for 3 days
- Children for 7 days
- Pregnant women for 15 days

**Distancing does not mean complete isolation. One way you can plan to keep in touch with friends and family is through phone or video calls.**



#### Wait for 7 days to have sex, then use birth control

During your entire course of treatment and for 14 weeks after your last dose, use effective birth control to prevent pregnancy.

### IMPORTANT SAFETY INFORMATION (continued)

#### What is the most important information I should know about PLUVICTO? (continued)

To minimize radiation exposure to others following administration of PLUVICTO, limit close contact (less than 3 feet) with household contacts for 2 days or with children and pregnant women for 7 days. Refrain from sexual activity for 7 days, and sleep in a separate bedroom from household contacts for 3 days, from children for 7 days, or from pregnant women for 15 days.



# Important reminders and tips after each dose of PLUVICTO

 *Print this page as a reminder for yourself and others.*

By following a few simple precautions in the first few days after treatment, you can reduce radiation exposure to others.

| Day 0<br>PLUVICTO<br>dose | Day 1<br>The day after<br>the dose | Day 2 | Day 3 | Days 4 to 7 | Days 8 to 15 |
|---------------------------|------------------------------------|-------|-------|-------------|--------------|
|---------------------------|------------------------------------|-------|-------|-------------|--------------|

**Stay more than 3 feet away, about an arm's length, from:**

|                                   |   |   |   |   |  |
|-----------------------------------|---|---|---|---|--|
| Adults                            | X | X |   |   |  |
| Children, pregnant women, or pets | X | X | X | X |  |

**Sleep alone in a separate room, away from:**

|                 |   |   |   |   |   |
|-----------------|---|---|---|---|---|
| Adults and pets | X | X | X |   |   |
| Children        | X | X | X | X |   |
| Pregnant women  | X | X | X | X | X |

|           |   |   |   |   |  |
|-----------|---|---|---|---|--|
| Avoid sex | X | X | X | X |  |
|-----------|---|---|---|---|--|

**Use effective birth control to prevent pregnancy during the entire course of treatment and for 14 weeks after your last dose**

**On the day of treatment:**



Help eliminate radiation from your body through frequent urination. Your doctor may recommend drinking about 8 ounces of fluid every hour on the day of your dose.

**On the way home from treatment:**



- If taking a car, sit behind an empty passenger seat so that you are safely distanced from the driver
- Avoid using public transportation for more than 1 hour during the first 3 days after treatment. When using it, do your best to follow the 3-foot distancing recommendation



## **At home:**

Because your body gives off radiation after each dose of PLUVICTO, especially in urine, follow the below practices for at least 2 days afterward to minimize radiation exposure to others.

Remember to drink plenty of water during those 2 days to help with frequent urination, and try to defecate (poop) every day.

- Consider sitting down to urinate to reduce splatter in the restroom
- Flush toilets twice and wash hands thoroughly
- Shower daily for at least the first 7 days after treatment
- Do not share towels and washcloths with others
- Wash your dishes separately from others
- Wash clothes that contain your sweat, blood, or urine separately from those of others when doing laundry
- Keep your plastic trash bags separate from other trash and away from children and animals



## **In public:**

- You will be given a Discharge Card. Carry it with you throughout treatment, especially during travel or hospital visits. Small amounts of radiation from your body may trigger radiation monitors
- Tell health care providers that you were treated with PLUVICTO if:
  - You need medical care during the first 2 weeks after a PLUVICTO dose, as there may still be some radiation in your body
  - You need to give blood, urine, or stool samples during the first 10 days after a PLUVICTO dose, as the samples might have radiation

**Ask your health care team for further advice and share this information  
with others as needed to help ensure their safety.**



[Click here to locate a treatment center near you.](#)



If you need help, ask your doctor or call Novartis Patient Support at 1-844-638-7222 to speak with a Novartis Patient Navigator.

**Once you connect with the treatment center, they do the rest, including confirming your insurance coverage.**

## IMPORTANT SAFETY INFORMATION (continued)

**PLUVICTO may cause serious side effects, including:**

**Low level of blood cell counts, which may be severe and life-threatening.** Tell your doctor right away if you develop any new or worsening symptoms, including:

- Tiredness or weakness
- Pale skin
- Shortness of breath
- Bleeding or bruising more easily than normal or difficulty stopping bleeding
- Frequent infections with signs such as fever, chills, sore throat, or mouth ulcers

**Please see additional Important Safety Information throughout.**

**Please see full Prescribing Information.**

**Novartis Patient Support is a comprehensive program designed to assist you every step of the way**

We can help you:



### 1:1 patient navigation support

A dedicated team is here to support and guide you through the referral process, financial assistance, and what to anticipate with treatment, and provide answers to common questions.



### Navigate the insurance process

The Novartis Patient Support team can work with your provider to help navigate insurance coverage for your medication.



### Get financial support\*

We offer savings options and resources to help you access treatment. If you have private insurance, you could be eligible for co-pay support and may pay as little as \$0 for your PLUVICTO treatment.

**[Click here to enroll or call us at 1-844-638-7222, Monday-Friday, 8:00 AM–8:00 PM ET, to speak to a Novartis Patient Navigator.](#)**

\*Limitations apply. Valid only for those with private insurance. The Program provides that an eligible patient will be responsible for the first \$0 and then may receive assistance for up to a maximum of \$15,000 over the course of the treatment to cover eligible out-of-pocket costs for the product. The patient is responsible for any costs once the limit is reached. Program not valid (i) under Medicare, Medicaid, TRICARE, VA, DoD, or any other federal or state healthcare program, (ii) where patient is not using insurance coverage at all, (iii) where the patient's insurance plan reimburses for the entire cost of the drug, or (iv) where product is not covered by patient's insurance. The value of this program is exclusively for the benefit of patients and is intended to be credited towards patient out-of-pocket obligations and maximums, including applicable co-payments, coinsurance, and deductibles. Program is not valid where prohibited by law. Patient may not seek reimbursement for the value received from this program from other parties, including any health insurance program or plan, flexible spending account, or healthcare savings account. Patient is responsible for complying with any applicable limitations and requirements of their health plan related to the use of the Program. Valid only in the United States and Puerto Rico. This Program is not health insurance. Program may not be combined with any third-party rebate, coupon, or offer. Proof of purchase may be required. Novartis reserves the right to rescind, revoke, or amend the Program and discontinue support at any time without notice.

A photograph of an older man with a beard and glasses, wearing a plaid shirt, standing behind a child who is wearing a green helmet and sitting in a blue go-kart. A small dog is also in the kart. The background is a large, stylized 'V' shape in blue and orange.

 **PLUVICTO<sup>®</sup>**  
lutetium Lu 177 vipivotide tetraxetan  
INJECTION FOR INTRAVENOUS USE

**You've got this,  
but you also  
don't have to  
do it alone.**

Patient portrayal.

**Novartis Patient Support  
for 1-on-1 support every  
step of the way.  
Call 1-844-638-7222.**

**[Click here](#) to watch the stories  
of other men who have taken  
PLUVICTO, and their loved ones.**



**Novartis Pharmaceuticals Corporation**  
East Hanover, New Jersey 07936-1080