

For adults with active psoriatic arthritis (PsA), non-radiographic axial spondyloarthritis (nr-axSpA), or ankylosing spondylitis (AS).



MOVE AND FEEL BETTER WITH COSENTYX® THROUGH IV INFUSION.

**REAL
PATIENT**

CAROLYN

COSENTYX® is available in both IV infusion and self-injection dosing options. Carolyn takes COSENTYX through IV infusion and has been compensated for her time. All other patients appearing in this brochure take COSENTYX with self-injections and have been compensated for their time. Individual results may vary.

INDICATIONS

COSENTYX® is a prescription medicine used to treat:

- adults with active psoriatic arthritis (PsA)
- adults with active ankylosing spondylitis (AS)
- adults with active non-radiographic axial spondyloarthritis (nr-axSpA) and objective signs of inflammation

IMPORTANT SAFETY INFORMATION

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

Please see Important Safety Information throughout the brochure.
Please see full Prescribing Information, including Medication Guide.

*“I like that I’m getting
the same results from
the medication in a
different form.”*

—Carolyn



REAL
PATIENT

CAROLYN

Table of Contents

About COSENTYX®	4
IV infusion	6
Psoriatic arthritis	8
Results for psoriatic arthritis	10
nr-axSpA and AS	12
Results for nr-axSpA and AS	14
How COSENTYX works	18
Savings and support	20

IMPORTANT SAFETY INFORMATION (continued)

What is the most important information I should know about COSENTYX?

COSENTYX is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

Infections

COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have had serious infections while taking COSENTYX, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses. Some people have died from these infections.

- Your doctor should check you for TB before starting treatment with COSENTYX.
- If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with COSENTYX and during treatment with COSENTYX.
- Your doctor should watch you closely for signs and symptoms of TB during treatment with COSENTYX.
Do not use COSENTYX if you have an active TB infection.

EXPERIENCE *matters*

COSENTYX® is a biologic treatment proven to help treat the symptoms of psoriatic arthritis (PsA), non-radiographic axial spondyloarthritis (nr-axSpA), and ankylosing spondylitis (AS).

#1

COSENTYX is the #1 prescribed treatment of its kind.*

6M+

Over 6 million prescriptions have been filled in the United States.*

18
YEARS

Studied for over 18 years in 90 clinical trials.*

10+
YEARS

10 years of FDA approval.†



COSENTYX through IV infusion contains the same medicine and was shown to be as effective and safe as self-injection.‡§

*Across all indications combined. As of October 2025.

†Plaque psoriasis–Jan. 2015. Psoriatic arthritis–Jan. 2016.

‡IV infusion contains the same medicine and was shown to be as effective and safe as COSENTYX self-injection for adult patients with active PsA, AS, or nr-axSpA.

§The FDA approval of the intravenous (IV) administration (injected into a vein) of COSENTYX is based on data showing that the amount of medication in your body after it is given directly through IV was within the range that would be seen when given as subcutaneous (SC) administration (injection under the skin).



*"I feel great, my joints feel great.
I can grab a wrench and work on
my motorcycle."*

REAL
PATIENT

GARY

IMPORTANT SAFETY INFORMATION (continued)

Before starting COSENTYX, tell your doctor if you:

- are being treated for an infection
- have an infection that does not go away or that keeps coming back
- have TB or have been in close contact with someone with TB
- have or have had Hepatitis B
- think you have an infection or have symptoms of an infection such as: fevers, sweats, or chills; muscle aches; cough; shortness of breath; blood in your phlegm; weight loss; warm, red, or painful skin or sores on your body; diarrhea or stomach pain; burning when you urinate or urinate more often than normal

the **FIRST** and only **IV INFUSION** of its kind

What to expect



- IV infusions are administered every 4 weeks and take about 30 minutes
- You will be given COSENTYX® by a healthcare provider through a needle placed in your vein (infusion).
- Your healthcare provider determines your dose based on your weight.
- Each infusion is given by a healthcare provider at an infusion center or doctor's office, so you have support during the process.



Watch Carolyn's story

"The infusions take 30 minutes. I'll often read a book while I'm there."

REAL
PATIENT

CAROLYN

How to prepare



Before arriving for your infusion, you may be advised to be **well hydrated and dress comfortably**.



During your infusion, your healthcare provider may **check your vital signs, ask you some questions, and answer any questions you may have**.

Pro tip: Having a routine can make the infusion more comfortable. During the process, you will be seated, so you can listen to music, read a book, or scroll through social media.



Once your infusion is complete, schedule your next infusion. **If you miss an appointment to receive COSENTYX, make another appointment as soon as possible.**

IMPORTANT SAFETY INFORMATION (continued)

After starting COSENTYX, call your doctor right away if you have any signs of infection listed. Do not use COSENTYX if you have any signs of infection unless you are instructed to by your doctor.

What are the possible side effects of COSENTYX?

COSENTYX may cause serious side effects, including:

Serious allergic reactions

Serious allergic reactions can occur. Get emergency medical help right away if you get any of the following symptoms: feeling faint; swelling of your face, eyelids, lips, mouth, tongue, or throat; trouble breathing or throat tightness; chest tightness; skin rash or hives (red, itchy bumps).

If you have a severe allergic reaction, do not give another injection of COSENTYX.

Could you be at **RISK** for permanent joint damage?

Psoriatic arthritis is an inflammatory, autoimmune condition that can affect both the joints and skin. It isn't caused from overuse of joints or aging. It's the result of an overactive immune system. Symptoms can include painful and tender joints, swollen fingers and toes, and sometimes even scaly, itchy skin. **If left untreated, psoriatic arthritis can cause permanent joint damage.**

COSENTYX[®] helps stop further joint damage

In a clinical trial of patients with psoriatic arthritis, about 60% of patients taking COSENTYX 150 mg or 300 mg saw at least a 20% improvement in psoriatic arthritis symptoms at 16 weeks.*

Relief that can last

84%

of people with psoriatic arthritis taking COSENTYX experienced no further joint damage at the 2-year mark.^{†‡}

*Vs 18% of patients taking placebo.

[†]Everyone's experience is different. These results were from a less rigorous analysis, so it is not known if symptom improvement was due to COSENTYX.

[‡]About 76% of patients taking COSENTYX 150 mg or 300 mg had no further joint damage at the hands, wrists, and feet at Week 24, compared to 68% taking placebo. At 2 years, 84% of patients taking COSENTYX 300 mg had no further joint damage (71% of patients taking COSENTYX 150 mg).

Please see Important Safety Information throughout the brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#). 8



"Real relief means movement, to be a dad, to be a grandpa, just to do life."

REAL
PATIENT

JORGE

IMPORTANT SAFETY INFORMATION (continued)

Inflammatory bowel disease

New cases of inflammatory bowel disease or "flare-ups" can happen with COSENTYX and can sometimes be serious. If you have inflammatory bowel disease (ulcerative colitis or Crohn's disease), tell your doctor if you have worsening disease symptoms during treatment with COSENTYX or develop new symptoms of stomach pain or diarrhea.


Severe skin reactions that look like eczema can happen during treatment with COSENTYX from days to months after your first dose and can sometimes lead to hospitalization. Your doctor may temporarily stop treatment with COSENTYX if you develop severe skin reactions. Tell your doctor if you have any of the following signs or symptoms: redness or rash; itching; small bumps or patches; your skin is dry or feels like leather; blisters on the hands or feet that ooze or become crusty or skin peeling.

 **Cosentyx[®]**
(secukinumab)

COSENTYX®

works fast for results that can last.

COSENTYX treats the multiple symptoms of psoriatic arthritis.



*In a clinical trial of adults taking 300 mg at 16 weeks.

†In a clinical trial of adults taking COSENTYX 300 mg as their first biologic at 12 weeks.

Please see Important Safety Information throughout the brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#). 10

Watch Elizabeth's Story



REAL
PATIENT

ELIZABETH

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of COSENTYX include:
cold symptoms, diarrhea, and upper respiratory tract infections.

These are not all of the possible side effects of COSENTYX. Call your doctor for medical advice about side effects.

 **Cosentyx®**
(secukinumab)

what is **AXIAL SPONDYLOARTHRITIS (axSpA)**?

axSpA is a painful inflammatory disease that can take a serious toll on your day-to-day. Living with back pain and morning stiffness can be painful. axSpA includes two underdiagnosed conditions:

- **Non-radiographic axial spondyloarthritis (nr-axSpA)**
- **Ankylosing spondylitis (AS)**

In both conditions, inflammation in the spinal joints can cause chronic back pain, stiffness, fatigue, and reduced mobility. **If left untreated, this inflammation can cause permanent damage.**

IMPORTANT SAFETY INFORMATION (continued)

Before using COSENTYX, tell your doctor if you:

- have any of the conditions or symptoms listed for infections.
- have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- are allergic to latex. The removable needle cap on the COSENTYX Sensoready[®] pen and prefilled syringes (150 mg/mL, 75 mg/0.5 mL) contains latex.

"I'm feeling better, I'm moving better, I'm able to get out and do the things that I want to do."



**REAL
PATIENT**

WENDY

nr-axSpA and AS

RESULTS *fast.* RELIEF *that can last.*

Many people taking COSENTYX® saw relief from symptoms in as little as 16 weeks, with similar results at 1 year.

nr-axSpA

40% Overall symptom improvement*^{†‡}

16 Weeks

1 Year

AS

3X Improvement in mobility^{†§||}

2X More back pain relief^{†¶}

16 Weeks

1 Year

5 YRS Many continued to feel relief at 5 years^{†#}

16 Weeks

5 Years

*41% of people on COSENTYX 150 mg at Week 16 vs 28% taking placebo.

†38% of people on COSENTYX 150 mg at 1 year vs 19% taking placebo.

‡All results shown in clinical trials using subcutaneous (SC) administration (injection under the skin).

§Reported by those in a clinical trial taking COSENTYX and who completed a questionnaire that asked how difficult it was to perform physical activities.

||At 16 weeks, patients saw 35% improvement in mobility while taking COSENTYX vs 11% taking placebo.

¶At 16 weeks, patients taking COSENTYX saw 43% improvement in back pain vs 16% taking placebo.

#20% improvement in overall symptoms vs those taking placebo (61% vs 28%).

Please see Important Safety Information throughout the brochure.
Please see full Prescribing Information, including Medication Guide. 14



REAL
PATIENT

MARK

Results for
nr-axSpA and AS

IMPORTANT SAFETY INFORMATION (continued)

•have recently received or are scheduled to receive an immunization (vaccine). People who take COSENTYX **should not** receive live vaccines. Children should be brought up to date with all vaccines before starting COSENTYX.

 **Cosentyx**[®]
(secukinumab)

MOVE *and* FEEL BETTER

When people taking COSENTYX® were given a survey by their doctors that measured how well their treatment was working,* the overall results showed improvements in some or all of the following symptoms of AS and nr-axSpA.



Total spinal pain



Morning stiffness



Fatigue



Tenderness (enthesitis)



Joint pain and swelling

*Bath Ankylosing Spondylitis Disease Activity Index.

Most people who took COSENTYX not only had less back pain and morning stiffness, they also saw improvement in their ability to move, and felt better overall.

IMPORTANT SAFETY INFORMATION (continued)

- have any other medical conditions and all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of your medicines to show your healthcare provider and pharmacist when you get a new medicine.

Watch Cat's Story



*"Before COSENTYX,
I couldn't do an essential stretch."*

REAL
PATIENT

CAT

IMPORTANT SAFETY INFORMATION (continued)


- are pregnant or plan to become pregnant. It is not known if COSENTYX can harm your unborn baby. You and your doctor should decide if you will use COSENTYX.
- are breastfeeding or plan to breastfeed. It is not known if COSENTYX passes into your breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

how does **COSENTYX®** work?

COSENTYX works differently by targeting and blocking IL-17A, a molecule in the body believed to play a role in inflammation in people with PsA, nr-axSpA, and AS.*

*The relationship between how COSENTYX works in the body and how it affects your symptoms is unknown.



Patient portrayal of how COSENTYX works.



Select your indication to see how **COSENTYX** works

Explore PsO

Explore PsA

Explore AS

Explore nr-axSpA

IMPORTANT SAFETY INFORMATION

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

What is the most important information I should know about COSENTYX?

COSENTYX is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

Infections

COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have had serious infections while taking COSENTYX, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses. Some people have died from these infections.

COST SHOULDN'T BE A BARRIER *between you and your treatment*

Novartis Patient Support will always try to help

Novartis Patient Support is a comprehensive program designed to help you start, stay, and save on COSENTYX®.



Novartis Patient Support

- Finding answers to questions
- Navigating insurance approvals and investigations
- Directing you to injection resources so you can start with confidence



Financial Support

If you can't afford your medicine, Novartis may be able to help.



Ongoing Support

The Novartis Patient Support team can offer additional helpful resources.

\$0 Co-Pay Plus offer

The Novartis Patient Support \$0 Co-Pay Plus* Offer

If you are privately insured, you may **pay as little as \$0 on COSENTYX** with Co-Pay Plus. The Co-Pay Plus offer can help pay for out-of-pocket costs related to your prescription, including refills. See Program Terms & Conditions for eligibility, or visit www.cosentyx.com/cost. You can also reach out directly to Novartis Patient Support at 844-COSENTYX (844-267-3689) Monday-Friday, 8:00 AM-8:00 PM ET, excluding holidays.

Please see Important Safety Information throughout the brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#). 20

"Novartis Patient Support has been nothing but excellent in helping me. They make sure my insurance is set and my prescription is delivered on time."



**See How Novartis Patient
Support Can Help**

***Limitations apply.** Subject to annual co-pay benefit limit. Offer not valid under Medicare, Medicaid, or any other federal or state programs. Novartis reserves the right to rescind, revoke, or amend this program without notice. Additional limitations may apply. See complete Terms & Conditions at cosentyx.com for details.

Coverage option

COSENTYX® IV infusions may be covered under Medicare Part B for people with PsA, AS, and nr-axSpA. Be sure to check with your insurance provider and talk to your doctor to understand your coverage and treatment options.

REAL
PATIENT

CAROLYN



"You don't have to self-administer the medication. IV infusion is a great alternative for me."

REAL
PATIENT

CAROLYN

Ask your rheumatologist if COSENTYX[®] is right for you.



Learn more at [COSENTYX.com](https://www.cosentyx.com)



[@COSENTYX \(secukinumab\)](https://www.facebook.com/cosentyxsecukinumab)



[@COSENTYX \(secukinumab\)](https://www.instagram.com/cosentyxsecukinumab)

Please see Important Safety Information throughout the brochure.
Please see full [Prescribing Information](#), including [Medication Guide](#).

 NOVARTIS

Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936-1080