

What do I need to know AFTER EACH PLUVICTO® DOSE?

1. Help reduce radiation exposure to yourself and others

Your body, blood, and urine give off radiation for a while after getting PLUVICTO. So, it is important to follow these instructions. For all these instructions, count Day 1 as the day after you get PLUVICTO.



Stay hydrated

For 1 day drink a lot and urinate (pee) as much as you can



Sleep alone in a separate bedroom

For 3 days away from other **adults**

For 7 days away from **children**

For 15 days away from **pregnant women**



Stay more than 3 feet away from others

For 2 days from other **adults**

For 7 days from **children** or **pregnant women**



No sex at first, and then use effective birth control

For 7 days no sex

During your course of treatment and for 14 weeks after your last dose use effective birth control to avoid pregnancy

Treatment may cause infertility

2. Be on the lookout for side effects during and after your treatment

Tell your doctor about any side effects you have as they may be able to help.

The most common side effects of PLUVICTO are:

- Decreased blood cell counts
- Laboratory abnormalities
- Tiredness
- Dry mouth
- Nausea
- Kidney problems

3. Plan your next blood-work appointment with your doctor

You will need to have your blood work done again 1-2 weeks before the next time you get PLUVICTO.

For more information, please
visit [PLUVICTO.com](https://www.pluvicto.com)
or contact your doctor.

Please see Important Safety Information on the next page and full Prescribing Information.

 **PLUVICTO®**
lutetium Lu 177 vipivotide tetraxetan
INJECTION FOR INTRAVENOUS USE

IMPORTANT SAFETY INFORMATION

What is PLUVICTO® (lutetium Lu 177 vipivotide tetraxetan)?

PLUVICTO is a prescription treatment used to treat adults with metastatic prostate-specific membrane antigen-positive (PSMA+) prostate cancer and are:

- androgen pathway modulation-naïve or -sensitive (mAPMN/S) in combination with hormone therapy, or
- androgen pathway modulation-resistant (mAPMR) after hormone therapy, and
 - appropriate to delay chemotherapy, or
 - have received prior chemotherapy

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about PLUVICTO?

Use of PLUVICTO involves exposure to radioactivity. Long-term, accruing radiation exposure is associated with an increased risk for cancer. Drink plenty of water and urinate as often as possible during the first hours after administration.

To minimize radiation exposure to others following administration of PLUVICTO, limit close contact (less than 3 feet) with household contacts for 2 days or with children and pregnant women for 7 days. Refrain from sexual activity for 7 days, and sleep in a separate bedroom from household contacts for 3 days, from children for 7 days, or from pregnant women for 15 days.

PLUVICTO may cause serious side effects, including:

Low level of blood cell counts, which may be severe and life-threatening. Tell your doctor right away if you develop any new or worsening symptoms, including:

- Tiredness or weakness
- Pale skin
- Shortness of breath

- Bleeding or bruising more easily than normal or difficulty stopping bleeding
- Frequent infections with signs such as fever, chills, sore throat, or mouth ulcers

Kidney problems, which can be severe. You should stay well hydrated before and after treatment. Tell your doctor right away if you develop any new or worsening urinary symptoms.

All radiopharmaceuticals, including PLUVICTO, have the potential to **cause harm to an unborn baby.**

- You should use effective contraception during treatment with PLUVICTO and for 14 weeks after your last dose

PLUVICTO may cause temporary or permanent **infertility.**

The most common side effects of PLUVICTO include:

- Decreased blood cell counts
- Laboratory abnormalities
- Tiredness
- Dry mouth
- Nausea
- Kidney problems

These are not all of the possible side effects of PLUVICTO. Call your doctor for advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see full [Prescribing Information](#).